



CITY UNIVERSITY
LONDON

Introduction to PT-quantum mechanics, deformations of integrable models

Andreas Fring

Banaras Hindu University, Varanasi, May 2018



CITY UNIVERSITY
LONDON

Introduction to PT-quantum mechanics, deformations of integrable models

Andreas Fring

Banaras Hindu University, Varanasi, May 2018

based on: A. Fring,
PT-Symmetric Deformations of Nonlinear Integrable Systems,
chapter 9 in "PT Symmetry: An Introduction",
World Scientific Publishing Co., Singapore, 2018

Outline

- 1 Introduction to PT-quantum mechanics
- 2 PT-deformed quantum spin chains
- 3 PT-deformed Calogero-Moser-Sutherland models
- 4 PT-deformed KdV/Ito systems
- 5 Conclusions



Ubiquitous non-Hermitian Hamiltonians (examples from the literature)

- with the standard inner product X is not Hermitian

$$X^\dagger = X + 2\tau i\hbar P \quad \text{and} \quad P^\dagger = P$$

Supersymmetry (Darboux transformation)

Decompose Hamiltonian \mathcal{H} as:

$$\mathcal{H} = H_+ \oplus H_- = Q\tilde{Q} \oplus \tilde{Q}Q$$

Supersymmetry (Darboux transformation)

Decompose Hamiltonian \mathcal{H} as:

$$\mathcal{H} = H_+ \oplus H_- = Q\tilde{Q} \oplus \tilde{Q}Q$$

- intertwining operators: $QH_- = H_+Q$ and $\tilde{Q}H_+ = H_-\tilde{Q}$

$$\Rightarrow [\mathcal{H}, Q] = [\mathcal{H}, \tilde{Q}] = 0$$

Supersymmetry (Darboux transformation)

Decompose Hamiltonian \mathcal{H} as:

$$\mathcal{H} = H_+ \oplus H_- = Q\tilde{Q} \oplus \tilde{Q}Q$$

- intertwining operators: $QH_- = H_+Q$ and $\tilde{Q}H_+ = H_-\tilde{Q}$

$$\Rightarrow [\mathcal{H}, Q] = [\mathcal{H}, \tilde{Q}] = 0$$

- realization: $Q = \frac{d}{dx} + W$ and $\tilde{Q} = -\frac{d}{dx} + W$

$$\Rightarrow H_{\pm} = -\Delta + W^2 \pm W' = -\Delta + V_{\pm}$$

Supersymmetry (Darboux transformation)

Decompose Hamiltonian \mathcal{H} as:

$$\mathcal{H} = H_+ \oplus H_- = Q\tilde{Q} \oplus \tilde{Q}Q$$

- intertwining operators: $QH_- = H_+Q$ and $\tilde{Q}H_+ = H_-\tilde{Q}$

$$\Rightarrow [\mathcal{H}, Q] = [\mathcal{H}, \tilde{Q}] = 0$$

- realization: $Q = \frac{d}{dx} + W$ and $\tilde{Q} = -\frac{d}{dx} + W$

$$\Rightarrow H_{\pm} = -\Delta + W^2 \pm W' = -\Delta + V_{\pm}$$

- ground state: $H_- \Phi_n^- = \varepsilon_n \Phi_n^-$ and $H_- \Phi_m^- = 0$

Supersymmetry (Darboux transformation)

Decompose Hamiltonian \mathcal{H} as:

$$\mathcal{H} = H_+ \oplus H_- = Q\tilde{Q} \oplus \tilde{Q}Q$$

- intertwining operators: $QH_- = H_+Q$ and $\tilde{Q}H_+ = H_-\tilde{Q}$

$$\Rightarrow [\mathcal{H}, Q] = [\mathcal{H}, \tilde{Q}] = 0$$

- realization: $Q = \frac{d}{dx} + W$ and $\tilde{Q} = -\frac{d}{dx} + W$

$$\Rightarrow H_{\pm} = -\Delta + W^2 \pm W' = -\Delta + V_{\pm}$$

- ground state: $H_- \Phi_n^- = \varepsilon_n \Phi_n^-$ and $H_- \Phi_m^- = 0$
 \Rightarrow isospectral Hamiltonians

$$H_{\pm}^m = -\Delta + V_{\pm}^m + E_m \quad H_{\pm}^m \Phi_n^{\pm} = E_n \Phi_n^{\pm} \quad \text{for } n > m$$

Unbroken \mathcal{PT} -symmetry guarantees real eigenvalues (QM)

- \mathcal{PT} -symmetry: $\mathcal{PT} : x \rightarrow -x \quad p \rightarrow p \quad i \rightarrow -i$
($\mathcal{P} : x \rightarrow -x, p \rightarrow -p$; $\mathcal{T} : x \rightarrow x, p \rightarrow -p, i \rightarrow -i$)

Unbroken \mathcal{PT} -symmetry guarantees real eigenvalues (QM)

- \mathcal{PT} -symmetry: $\mathcal{PT} : x \rightarrow -x \quad p \rightarrow p \quad i \rightarrow -i$
 $(\mathcal{P} : x \rightarrow -x, p \rightarrow -p; \mathcal{T} : x \rightarrow x, p \rightarrow -p, i \rightarrow -i)$
- \mathcal{PT} is an anti-linear operator:

$$\mathcal{PT}(\lambda\Phi + \mu\Psi) = \lambda^*\mathcal{PT}\Phi + \mu^*\mathcal{PT}\Psi \quad \lambda, \mu \in \mathbb{C}$$

Unbroken \mathcal{PT} -symmetry guarantees real eigenvalues (QM)

- \mathcal{PT} -symmetry: $\mathcal{PT} : x \rightarrow -x \quad p \rightarrow p \quad i \rightarrow -i$
 $(\mathcal{P} : x \rightarrow -x, p \rightarrow -p; \mathcal{T} : x \rightarrow x, p \rightarrow -p, i \rightarrow -i)$
- \mathcal{PT} is an anti-linear operator:

$$\mathcal{PT}(\lambda\Phi + \mu\Psi) = \lambda^*\mathcal{PT}\Phi + \mu^*\mathcal{PT}\Psi \quad \lambda, \mu \in \mathbb{C}$$

- Real eigenvalues from unbroken \mathcal{PT} -symmetry:

$$[\mathcal{H}, \mathcal{PT}] = 0 \quad \wedge \quad \mathcal{PT}\Phi = \Phi \quad \Rightarrow \quad \varepsilon = \varepsilon^* \quad \text{for } \mathcal{H}\Phi = \varepsilon\Phi$$

Unbroken \mathcal{PT} -symmetry guarantees real eigenvalues (QM)

- \mathcal{PT} -symmetry: $\mathcal{PT} : x \rightarrow -x \quad p \rightarrow p \quad i \rightarrow -i$
 $(\mathcal{P} : x \rightarrow -x, p \rightarrow -p; \mathcal{T} : x \rightarrow x, p \rightarrow -p, i \rightarrow -i)$
- \mathcal{PT} is an anti-linear operator:

$$\mathcal{PT}(\lambda\Phi + \mu\Psi) = \lambda^*\mathcal{PT}\Phi + \mu^*\mathcal{PT}\Psi \quad \lambda, \mu \in \mathbb{C}$$

- Real eigenvalues from unbroken \mathcal{PT} -symmetry:

$$[\mathcal{H}, \mathcal{PT}] = 0 \quad \wedge \quad \mathcal{PT}\Phi = \Phi \quad \Rightarrow \quad \varepsilon = \varepsilon^* \quad \text{for } \mathcal{H}\Phi = \varepsilon\Phi$$

- *Proof:*

Unbroken \mathcal{PT} -symmetry guarantees real eigenvalues (QM)

- \mathcal{PT} -symmetry: $\mathcal{PT} : x \rightarrow -x \quad p \rightarrow p \quad i \rightarrow -i$
 $(\mathcal{P} : x \rightarrow -x, p \rightarrow -p; \mathcal{T} : x \rightarrow x, p \rightarrow -p, i \rightarrow -i)$
- \mathcal{PT} is an anti-linear operator:

$$\mathcal{PT}(\lambda\Phi + \mu\Psi) = \lambda^*\mathcal{PT}\Phi + \mu^*\mathcal{PT}\Psi \quad \lambda, \mu \in \mathbb{C}$$

- Real eigenvalues from unbroken \mathcal{PT} -symmetry:

$$[\mathcal{H}, \mathcal{PT}] = 0 \quad \wedge \quad \mathcal{PT}\Phi = \Phi \quad \Rightarrow \quad \varepsilon = \varepsilon^* \quad \text{for } \mathcal{H}\Phi = \varepsilon\Phi$$

- *Proof:*

$$\varepsilon\Phi = \mathcal{H}\Phi$$

Unbroken \mathcal{PT} -symmetry guarantees real eigenvalues (QM)

- \mathcal{PT} -symmetry: $\mathcal{PT} : x \rightarrow -x \quad p \rightarrow p \quad i \rightarrow -i$
 $(\mathcal{P} : x \rightarrow -x, p \rightarrow -p; \mathcal{T} : x \rightarrow x, p \rightarrow -p, i \rightarrow -i)$
- \mathcal{PT} is an anti-linear operator:

$$\mathcal{PT}(\lambda\Phi + \mu\Psi) = \lambda^*\mathcal{PT}\Phi + \mu^*\mathcal{PT}\Psi \quad \lambda, \mu \in \mathbb{C}$$

- Real eigenvalues from unbroken \mathcal{PT} -symmetry:

$$[\mathcal{H}, \mathcal{PT}] = 0 \quad \wedge \quad \mathcal{PT}\Phi = \Phi \quad \Rightarrow \quad \varepsilon = \varepsilon^* \quad \text{for } \mathcal{H}\Phi = \varepsilon\Phi$$

- *Proof:*

$$\varepsilon\Phi = \mathcal{H}\Phi = \mathcal{H}\mathcal{PT}\Phi$$

Unbroken \mathcal{PT} -symmetry guarantees real eigenvalues (QM)

- \mathcal{PT} -symmetry: $\mathcal{PT} : x \rightarrow -x \quad p \rightarrow p \quad i \rightarrow -i$
 $(\mathcal{P} : x \rightarrow -x, p \rightarrow -p; \mathcal{T} : x \rightarrow x, p \rightarrow -p, i \rightarrow -i)$
- \mathcal{PT} is an anti-linear operator:

$$\mathcal{PT}(\lambda\Phi + \mu\Psi) = \lambda^*\mathcal{PT}\Phi + \mu^*\mathcal{PT}\Psi \quad \lambda, \mu \in \mathbb{C}$$

- Real eigenvalues from unbroken \mathcal{PT} -symmetry:

$$[\mathcal{H}, \mathcal{PT}] = 0 \quad \wedge \quad \mathcal{PT}\Phi = \Phi \quad \Rightarrow \quad \varepsilon = \varepsilon^* \quad \text{for } \mathcal{H}\Phi = \varepsilon\Phi$$

- *Proof:*

$$\varepsilon\Phi = \mathcal{H}\Phi = \mathcal{H}\mathcal{PT}\Phi = \mathcal{PT}\mathcal{H}\Phi$$

Unbroken \mathcal{PT} -symmetry guarantees real eigenvalues (QM)

- \mathcal{PT} -symmetry: $\mathcal{PT} : x \rightarrow -x \quad p \rightarrow p \quad i \rightarrow -i$
 $(\mathcal{P} : x \rightarrow -x, p \rightarrow -p; \mathcal{T} : x \rightarrow x, p \rightarrow -p, i \rightarrow -i)$
- \mathcal{PT} is an anti-linear operator:

$$\mathcal{PT}(\lambda\Phi + \mu\Psi) = \lambda^*\mathcal{PT}\Phi + \mu^*\mathcal{PT}\Psi \quad \lambda, \mu \in \mathbb{C}$$

- Real eigenvalues from unbroken \mathcal{PT} -symmetry:

$$[\mathcal{H}, \mathcal{PT}] = 0 \quad \wedge \quad \mathcal{PT}\Phi = \Phi \quad \Rightarrow \quad \varepsilon = \varepsilon^* \quad \text{for } \mathcal{H}\Phi = \varepsilon\Phi$$

- *Proof:*

$$\varepsilon\Phi = \mathcal{H}\Phi = \mathcal{H}\mathcal{PT}\Phi = \mathcal{PT}\mathcal{H}\Phi = \mathcal{PT}\varepsilon\Phi$$

Unbroken \mathcal{PT} -symmetry guarantees real eigenvalues (QM)

- \mathcal{PT} -symmetry: $\mathcal{PT} : x \rightarrow -x \quad p \rightarrow p \quad i \rightarrow -i$
 $(\mathcal{P} : x \rightarrow -x, p \rightarrow -p; \mathcal{T} : x \rightarrow x, p \rightarrow -p, i \rightarrow -i)$
- \mathcal{PT} is an anti-linear operator:

$$\mathcal{PT}(\lambda\Phi + \mu\Psi) = \lambda^*\mathcal{PT}\Phi + \mu^*\mathcal{PT}\Psi \quad \lambda, \mu \in \mathbb{C}$$

- Real eigenvalues from unbroken \mathcal{PT} -symmetry:

$$[\mathcal{H}, \mathcal{PT}] = 0 \quad \wedge \quad \mathcal{PT}\Phi = \Phi \quad \Rightarrow \quad \varepsilon = \varepsilon^* \quad \text{for } \mathcal{H}\Phi = \varepsilon\Phi$$

- *Proof:*

$$\varepsilon\Phi = \mathcal{H}\Phi = \mathcal{H}\mathcal{PT}\Phi = \mathcal{PT}\mathcal{H}\Phi = \mathcal{PT}\varepsilon\Phi = \varepsilon^*\mathcal{PT}\Phi$$

Unbroken \mathcal{PT} -symmetry guarantees real eigenvalues (QM)

- \mathcal{PT} -symmetry: $\mathcal{PT} : x \rightarrow -x \quad p \rightarrow p \quad i \rightarrow -i$
 $(\mathcal{P} : x \rightarrow -x, p \rightarrow -p; \mathcal{T} : x \rightarrow x, p \rightarrow -p, i \rightarrow -i)$
- \mathcal{PT} is an anti-linear operator:

$$\mathcal{PT}(\lambda\Phi + \mu\Psi) = \lambda^*\mathcal{PT}\Phi + \mu^*\mathcal{PT}\Psi \quad \lambda, \mu \in \mathbb{C}$$

- Real eigenvalues from unbroken \mathcal{PT} -symmetry:

$$[\mathcal{H}, \mathcal{PT}] = 0 \quad \wedge \quad \mathcal{PT}\Phi = \Phi \quad \Rightarrow \quad \varepsilon = \varepsilon^* \quad \text{for } \mathcal{H}\Phi = \varepsilon\Phi$$

- *Proof:*

$$\varepsilon\Phi = \mathcal{H}\Phi = \mathcal{H}\mathcal{PT}\Phi = \mathcal{PT}\mathcal{H}\Phi = \mathcal{PT}\varepsilon\Phi = \varepsilon^*\mathcal{PT}\Phi = \varepsilon^*\Phi$$

Unbroken \mathcal{PT} -symmetry guarantees real eigenvalues (QM)

- \mathcal{PT} -symmetry: $\mathcal{PT} : x \rightarrow -x \quad p \rightarrow p \quad i \rightarrow -i$
 $(\mathcal{P} : x \rightarrow -x, p \rightarrow -p; \mathcal{T} : x \rightarrow x, p \rightarrow -p, i \rightarrow -i)$
- \mathcal{PT} is an anti-linear operator:

$$\mathcal{PT}(\lambda\Phi + \mu\Psi) = \lambda^*\mathcal{PT}\Phi + \mu^*\mathcal{PT}\Psi \quad \lambda, \mu \in \mathbb{C}$$

- Real eigenvalues from unbroken \mathcal{PT} -symmetry:

$$[\mathcal{H}, \mathcal{PT}] = 0 \quad \wedge \quad \mathcal{PT}\Phi = \Phi \quad \Rightarrow \quad \varepsilon = \varepsilon^* \quad \text{for } \mathcal{H}\Phi = \varepsilon\Phi$$

- *Proof:*

$$\varepsilon\Phi = \mathcal{H}\Phi = \mathcal{H}\mathcal{PT}\Phi = \mathcal{PT}\mathcal{H}\Phi = \mathcal{PT}\varepsilon\Phi = \varepsilon^*\mathcal{PT}\Phi = \varepsilon^*\Phi$$

PT-symmetry is only an example of an antilinear involution

How to formulate a quantum mechanical framework?

- 1 orthogonality
- 2 observables
- 3 uniqueness
- 4 technicalities (new metric etc)

Orthogonality

- Take h to be a Hermitian and diagonalisable Hamiltonian:

$$\langle \phi_n | h \phi_m \rangle = \langle h \phi_n | \phi_m \rangle$$

$$\begin{aligned} |h \phi_m \rangle &= \varepsilon_m |\phi_m \rangle \\ \langle h \phi_n | &= \varepsilon_n^* \langle \phi_n | \end{aligned}$$

Orthogonality

- Take h to be a Hermitian and diagonalisable Hamiltonian:

$$\langle \phi_n | h \phi_m \rangle = \langle h \phi_n | \phi_m \rangle$$

$$\langle \phi_n | h \phi_m \rangle = \varepsilon_m \langle \phi_n | \phi_m \rangle$$

$$\langle h \phi_n | \phi_m \rangle = \varepsilon_n^* \langle \phi_n | \phi_m \rangle$$

Orthogonality

- Take h to be a Hermitian and diagonalisable Hamiltonian:

$$\langle \phi_n | h \phi_m \rangle = \langle h \phi_n | \phi_m \rangle$$

$$\left. \begin{array}{l} \langle \phi_n | h \phi_m \rangle = \varepsilon_m \langle \phi_n | \phi_m \rangle \\ \langle h \phi_n | \phi_m \rangle = \varepsilon_n^* \langle \phi_n | \phi_m \rangle \end{array} \right\} \Rightarrow 0 = (\varepsilon_m - \varepsilon_n^*) \langle \phi_n | \phi_m \rangle$$

Orthogonality

- Take h to be a Hermitian and diagonalisable Hamiltonian:

$$\langle \phi_n | h \phi_m \rangle = \langle h \phi_n | \phi_m \rangle$$

$$\left. \begin{array}{l} \langle \phi_n | h \phi_m \rangle = \varepsilon_m \langle \phi_n | \phi_m \rangle \\ \langle h \phi_n | \phi_m \rangle = \varepsilon_n^* \langle \phi_n | \phi_m \rangle \end{array} \right\} \Rightarrow 0 = (\varepsilon_m - \varepsilon_n^*) \langle \phi_n | \phi_m \rangle$$

$$\Rightarrow \quad n = m : \varepsilon_n = \varepsilon_n^* \quad n \neq m : \langle \phi_n | \phi_m \rangle = 0$$

Orthogonality

- Take h to be a Hermitian and diagonalisable Hamiltonian:

$$\langle \phi_n | h \phi_m \rangle = \langle h \phi_n | \phi_m \rangle$$

$$\left. \begin{aligned} \langle \phi_n | h \phi_m \rangle &= \varepsilon_m \langle \phi_n | \phi_m \rangle \\ \langle h \phi_n | \phi_m \rangle &= \varepsilon_n^* \langle \phi_n | \phi_m \rangle \end{aligned} \right\} \Rightarrow 0 = (\varepsilon_m - \varepsilon_n^*) \langle \phi_n | \phi_m \rangle$$

$$\Rightarrow \quad n = m : \varepsilon_n = \varepsilon_n^* \quad n \neq m : \langle \phi_n | \phi_m \rangle = 0$$

- Take H to be a non-Hermitian Hamiltonian:

$$H |\Phi_n\rangle = \varepsilon_n |\Phi_n\rangle$$

- reality and orthogonality no longer guaranteed. Define

$$\langle \Phi_n | \Phi_m \rangle_\eta := \langle \Phi_n | \eta^2 \Phi_m \rangle$$

- when $\langle \Phi_n | H \Phi_m \rangle_\eta = \langle H \Phi_n | \Phi_m \rangle_\eta \Rightarrow \langle \Phi_n | \Phi_m \rangle_\eta = \delta_{n,m}$

H is Hermitian with respect to new metric

- Assume pseudo-Hermiticity:

$$h = \eta H \eta^{-1} = h^\dagger = (\eta^{-1})^\dagger H^\dagger \eta^\dagger \Leftrightarrow H^\dagger \eta^\dagger \eta = \eta^\dagger \eta H$$

$$\Phi = \eta^{-1} \phi \quad \eta^\dagger = \eta$$

$\Rightarrow H$ is Hermitian with respect to the new metric

H is Hermitian with respect to new metric

- Assume pseudo-Hermiticity:

$$h = \eta H \eta^{-1} = h^\dagger = (\eta^{-1})^\dagger H^\dagger \eta^\dagger \Leftrightarrow H^\dagger \eta^\dagger \eta = \eta^\dagger \eta H$$

$$\Phi = \eta^{-1} \phi \quad \eta^\dagger = \eta$$

$\Rightarrow H$ is Hermitian with respect to the new metric

Proof:

$$\langle \Psi | H \Phi \rangle_\eta = \langle \Psi | \eta^2 H \Phi \rangle = \langle \eta^{-1} \psi | \eta^2 H \eta^{-1} \phi \rangle$$

H is Hermitian with respect to new metric

- Assume pseudo-Hermiticity:

$$h = \eta H \eta^{-1} = h^\dagger = (\eta^{-1})^\dagger H^\dagger \eta^\dagger \Leftrightarrow H^\dagger \eta^\dagger \eta = \eta^\dagger \eta H$$

$$\Phi = \eta^{-1} \phi \quad \eta^\dagger = \eta$$

$\Rightarrow H$ is Hermitian with respect to the new metric

Proof:

$$\langle \Psi | H \Phi \rangle_\eta = \langle \Psi | \eta^2 H \Phi \rangle = \langle \eta^{-1} \psi | \eta^2 H \eta^{-1} \phi \rangle = \langle \psi | \eta H \eta^{-1} \phi \rangle =$$

$$\langle \psi | h \phi \rangle$$

H is Hermitian with respect to new metric

- Assume pseudo-Hermiticity:

$$h = \eta H \eta^{-1} = h^\dagger = (\eta^{-1})^\dagger H^\dagger \eta^\dagger \Leftrightarrow H^\dagger \eta^\dagger \eta = \eta^\dagger \eta H$$

$$\Phi = \eta^{-1} \phi \quad \eta^\dagger = \eta$$

$\Rightarrow H$ is Hermitian with respect to the new metric

Proof:

$$\begin{aligned} \langle \Psi | H \Phi \rangle_\eta &= \langle \Psi | \eta^2 H \Phi \rangle = \langle \eta^{-1} \psi | \eta^2 H \eta^{-1} \phi \rangle = \langle \psi | \eta H \eta^{-1} \phi \rangle = \\ \langle \psi | h \phi \rangle &= \langle h \psi | \phi \rangle = \langle \eta H \eta^{-1} \psi | \phi \rangle \end{aligned}$$

H is Hermitian with respect to new metric

- Assume pseudo-Hermiticity:

$$h = \eta H \eta^{-1} = h^\dagger = (\eta^{-1})^\dagger H^\dagger \eta^\dagger \Leftrightarrow H^\dagger \eta^\dagger \eta = \eta^\dagger \eta H$$

$$\Phi = \eta^{-1} \phi \quad \eta^\dagger = \eta$$

$\Rightarrow H$ is Hermitian with respect to the new metric

Proof:

$$\begin{aligned} \langle \Psi | H \Phi \rangle_\eta &= \langle \Psi | \eta^2 H \Phi \rangle = \langle \eta^{-1} \psi | \eta^2 H \eta^{-1} \phi \rangle = \langle \psi | \eta H \eta^{-1} \phi \rangle = \\ &\langle \psi | h \phi \rangle = \langle h \psi | \phi \rangle = \langle \eta H \eta^{-1} \psi | \phi \rangle = \langle H \Psi | \eta \phi \rangle = \langle H \Psi | \eta^2 \phi \rangle \\ &= \langle H \Psi | \Phi \rangle_\eta \end{aligned}$$

H is Hermitian with respect to new metric

- Assume pseudo-Hermiticity:

$$h = \eta H \eta^{-1} = h^\dagger = (\eta^{-1})^\dagger H^\dagger \eta^\dagger \Leftrightarrow H^\dagger \eta^\dagger \eta = \eta^\dagger \eta H$$

$$\Phi = \eta^{-1} \phi \quad \eta^\dagger = \eta$$

$\Rightarrow H$ is Hermitian with respect to the new metric

Proof:

$$\begin{aligned} \langle \Psi | H \Phi \rangle_\eta &= \langle \Psi | \eta^2 H \Phi \rangle = \langle \eta^{-1} \psi | \eta^2 H \eta^{-1} \phi \rangle = \langle \psi | \eta H \eta^{-1} \phi \rangle = \\ &= \langle \psi | h \phi \rangle = \langle h \psi | \phi \rangle = \langle \eta H \eta^{-1} \psi | \phi \rangle = \langle H \Psi | \eta \phi \rangle = \langle H \Psi | \eta^2 \phi \rangle \\ &= \langle H \Psi | \Phi \rangle_\eta \end{aligned}$$

\Rightarrow Eigenvalues of H are real, eigenstates are orthogonal

CPT-metric

[Bender, Brody, Jones, Phys. Rev. Lett. 89 (2002) 270401]

$$\langle \Psi | \Phi \rangle_{\mathcal{CPT}} := (\mathcal{CPT} | \Psi \rangle)^T \cdot | \Phi \rangle$$

- In position space: $\mathcal{C}(x, y) = \sum_n \Phi_n(x) \Phi_n(y)$
Very formal as normally one does not know $\Phi_n(x) \forall n$
- Algebraic approach: Solve
 $\mathcal{C}^2 = \mathbb{I} \quad [\mathcal{H}, \mathcal{C}] = 0 \quad [\mathcal{C}, \mathcal{PT}] = 0 \quad [\mathcal{H}, \mathcal{PT}] = 0$
- Relation \mathcal{C} and metric (same as pseudo-Hermiticity)

$$\mathcal{C} = \rho^{-1} \mathcal{P}$$

CPT-metric

[Bender, Brody, Jones, Phys. Rev. Lett. 89 (2002) 270401]

$$\langle \Psi | \Phi \rangle_{\mathcal{CPT}} := (\mathcal{CPT} | \Psi \rangle)^T \cdot | \Phi \rangle$$

- In position space: $\mathcal{C}(x, y) = \sum_n \Phi_n(x) \Phi_n(y)$
Very formal as normally one does not know $\Phi_n(x) \forall n$
- Algebraic approach: Solve
 $\mathcal{C}^2 = \mathbb{I} \quad [\mathcal{H}, \mathcal{C}] = 0 \quad [\mathcal{C}, \mathcal{PT}] = 0 \quad [\mathcal{H}, \mathcal{PT}] = 0$
- Relation \mathcal{C} and metric (same as pseudo-Hermiticity)

$$\mathcal{C} = \rho^{-1} \mathcal{P}$$

Proof:

CPT-metric

[Bender, Brody, Jones, Phys. Rev. Lett. 89 (2002) 270401]

$$\langle \Psi | \Phi \rangle_{\mathcal{CPT}} := (\mathcal{CPT} | \Psi \rangle)^T \cdot | \Phi \rangle$$

- In position space: $\mathcal{C}(x, y) = \sum_n \Phi_n(x) \Phi_n(y)$
Very formal as normally one does not know $\Phi_n(x) \forall n$
- Algebraic approach: Solve
 $\mathcal{C}^2 = \mathbb{I} \quad [\mathcal{H}, \mathcal{C}] = 0 \quad [\mathcal{C}, \mathcal{PT}] = 0 \quad [\mathcal{H}, \mathcal{PT}] = 0$
- Relation \mathcal{C} and metric (same as pseudo-Hermiticity)

$$\mathcal{C} = \rho^{-1} \mathcal{P}$$

Proof:

$$\rho \mathcal{H} = \mathcal{P} \mathcal{C} \mathcal{H}$$

CPT-metric

[Bender, Brody, Jones, Phys. Rev. Lett. 89 (2002) 270401]

$$\langle \Psi | \Phi \rangle_{\mathcal{CPT}} := (\mathcal{CPT} | \Psi \rangle)^T \cdot | \Phi \rangle$$

- In position space: $\mathcal{C}(x, y) = \sum_n \Phi_n(x) \Phi_n(y)$
Very formal as normally one does not know $\Phi_n(x) \forall n$
- Algebraic approach: Solve
 $\mathcal{C}^2 = \mathbb{I} \quad [\mathcal{H}, \mathcal{C}] = 0 \quad [\mathcal{C}, \mathcal{PT}] = 0 \quad [\mathcal{H}, \mathcal{PT}] = 0$
- Relation \mathcal{C} and metric (same as pseudo-Hermiticity)

$$\mathcal{C} = \rho^{-1} \mathcal{P}$$

Proof:

$$\rho \mathcal{H} = \mathcal{P} \mathcal{C} \mathcal{H} = \mathcal{P} \mathcal{H} \mathcal{C}$$

CPT-metric

[Bender, Brody, Jones, Phys. Rev. Lett. 89 (2002) 270401]

$$\langle \Psi | \Phi \rangle_{\mathcal{CPT}} := (\mathcal{CPT} | \Psi \rangle)^T \cdot | \Phi \rangle$$

- In position space: $\mathcal{C}(x, y) = \sum_n \Phi_n(x) \Phi_n(y)$
Very formal as normally one does not know $\Phi_n(x) \forall n$
- Algebraic approach: Solve
 $\mathcal{C}^2 = \mathbb{I} \quad [\mathcal{H}, \mathcal{C}] = 0 \quad [\mathcal{C}, \mathcal{PT}] = 0 \quad [\mathcal{H}, \mathcal{PT}] = 0$
- Relation \mathcal{C} and metric (same as pseudo-Hermiticity)

$$\mathcal{C} = \rho^{-1} \mathcal{P}$$

Proof: $\mathcal{P}^2 = \mathbb{I}$

$$\rho \mathcal{H} = \mathcal{P} \mathcal{C} \mathcal{H} = \mathcal{P} \mathcal{H} \mathcal{C} = \mathcal{P} \mathcal{P} \mathcal{T} \mathcal{H} \mathcal{P} \mathcal{T} \mathcal{C} = \mathcal{T} \mathcal{H} \mathcal{T} \mathcal{P} \mathcal{C}$$

CPT-metric

[Bender, Brody, Jones, Phys. Rev. Lett. 89 (2002) 270401]

$$\langle \Psi | \Phi \rangle_{\mathcal{CPT}} := (\mathcal{CPT} | \Psi \rangle)^T \cdot | \Phi \rangle$$

- In position space: $\mathcal{C}(x, y) = \sum_n \Phi_n(x) \Phi_n(y)$
Very formal as normally one does not know $\Phi_n(x) \forall n$
- Algebraic approach: Solve
 $\mathcal{C}^2 = \mathbb{I} \quad [\mathcal{H}, \mathcal{C}] = 0 \quad [\mathcal{C}, \mathcal{PT}] = 0 \quad [\mathcal{H}, \mathcal{PT}] = 0$
- Relation \mathcal{C} and metric (same as pseudo-Hermiticity)

$$\mathcal{C} = \rho^{-1} \mathcal{P}$$

Proof:

$$\rho \mathcal{H} = \mathcal{P} \mathcal{C} \mathcal{H} = \mathcal{P} \mathcal{H} \mathcal{C} = \mathcal{P} \mathcal{P} \mathcal{T} \mathcal{H} \mathcal{P} \mathcal{T} \mathcal{C} = \mathcal{T} \mathcal{H} \mathcal{T} \mathcal{P} \mathcal{C} = \mathcal{H}^\dagger \mathcal{P} \mathcal{C} = \mathcal{H}^\dagger \rho$$

CPT-metric

[Bender, Brody, Jones, Phys. Rev. Lett. 89 (2002) 270401]

$$\langle \Psi | \Phi \rangle_{\mathcal{CPT}} := (\mathcal{CPT} | \Psi \rangle)^T \cdot | \Phi \rangle$$

- In position space: $\mathcal{C}(x, y) = \sum_n \Phi_n(x) \Phi_n(y)$
Very formal as normally one does not know $\Phi_n(x) \forall n$
- Algebraic approach: Solve
 $\mathcal{C}^2 = \mathbb{I} \quad [\mathcal{H}, \mathcal{C}] = 0 \quad [\mathcal{C}, \mathcal{PT}] = 0 \quad [\mathcal{H}, \mathcal{PT}] = 0$
- Relation \mathcal{C} and metric (same as pseudo-Hermiticity)

$$\mathcal{C} = \rho^{-1} \mathcal{P}$$

Proof:

$$\rho \mathcal{H} = \mathcal{P} \mathcal{C} \mathcal{H} = \mathcal{P} \mathcal{H} \mathcal{C} = \mathcal{P} \mathcal{P} \mathcal{T} \mathcal{H} \mathcal{P} \mathcal{T} \mathcal{C} = \mathcal{T} \mathcal{H} \mathcal{T} \mathcal{P} \mathcal{C} = \mathcal{H}^\dagger \mathcal{P} \mathcal{C} = \mathcal{H}^\dagger \rho$$

Observables

- Observables are Hermitian with respect to the new metric

$$\langle \Phi_n | \mathcal{O} \Phi_m \rangle_\eta = \langle \mathcal{O} \Phi_n | \Phi_m \rangle_\eta$$

$$\mathcal{O} = \eta^{-1} o \eta \quad \Leftrightarrow \quad \mathcal{O}^\dagger = \rho \mathcal{O} \rho^{-1}$$

- o is an observable in the Hermitian system
- \mathcal{O} is an observable in the non-Hermitian system

General technique:

- Given H $\left\{ \begin{array}{l} \text{either solve } \eta H \eta^{-1} = h \text{ for } \eta \Rightarrow \rho = \eta^\dagger \eta \\ \text{or solve } H^\dagger = \rho H \rho^{-1} \text{ for } \rho \Rightarrow \eta = \sqrt{\rho} \end{array} \right.$

General technique:

- Given H $\left\{ \begin{array}{l} \text{either solve } \eta H \eta^{-1} = h \text{ for } \eta \Rightarrow \rho = \eta^\dagger \eta \\ \text{or solve } H^\dagger = \rho H \rho^{-1} \text{ for } \rho \Rightarrow \eta = \sqrt{\rho} \end{array} \right.$
- involves complicated commutation relations
- often this can only be solved perturbatively

Note:

- Thus, this is not re-inventing or disputing the validity of quantum mechanics.
- We only give up the restrictive requirement that Hamiltonians have to be Hermitian.

[C. Bender, *Rep. Prog. Phys.* 70 (2007) 947]

[A. Mostafazadeh, *Int. J. Geom. Meth. Phys.* 7 (2010) 1191]

[C. Bender, A. Fring et. al *PT-symmetric Quantum Mechanics*, Imperial College Press (2018?)]

Non-Hermitian time-dependent Hamiltonians

Is it possible to have a consistent description of time-dependent non-Hermitian Hamiltonian systems?

Non-Hermitian time-dependent Hamiltonians

Is it possible to have a consistent description of time-dependent non-Hermitian Hamiltonian systems?

Three scenarios

- time-dependent: $H(t) \neq H^\dagger(t)$, $h(t) = h^\dagger(t)$
time-independent: η, ρ

Non-Hermitian time-dependent Hamiltonians

Is it possible to have a consistent description of time-dependent non-Hermitian Hamiltonian systems?

Three scenarios

- time-dependent: $H(t) \neq H^\dagger(t)$, $h(t) = h^\dagger(t)$
time-independent: η, ρ
- time-dependent: $h(t) = h^\dagger(t)$, $\eta(t), \rho(t)$
time-independent: $H \neq H^\dagger$

Non-Hermitian time-dependent Hamiltonians

Is it possible to have a consistent description of time-dependent non-Hermitian Hamiltonian systems?

Three scenarios

- time-dependent: $H(t) \neq H^\dagger(t)$, $h(t) = h^\dagger(t)$
time-independent: η, ρ
- time-dependent: $h(t) = h^\dagger(t)$, $\eta(t), \rho(t)$
time-independent: $H \neq H^\dagger$
- time-dependent: $H(t) \neq H^\dagger(t)$, $h(t) = h^\dagger(t)$, $\eta(t), \rho(t)$

Theoretical framework (key equations):

Time-dependent Schrödinger eqn for $h(t) = h^\dagger(t)$, $H(t) \neq H^\dagger(t)$

$$h(t)\phi(t) = i\hbar\partial_t\phi(t), \quad \text{and} \quad H(t)\Psi(t) = i\hbar\partial_t\Psi(t)$$

Theoretical framework (key equations):

Time-dependent Schrödinger eqn for $h(t) = h^\dagger(t)$, $H(t) \neq H^\dagger(t)$

$$h(t)\phi(t) = i\hbar\partial_t\phi(t), \quad \text{and} \quad H(t)\Psi(t) = i\hbar\partial_t\Psi(t)$$

Time-dependent Dyson operator

$$\phi(t) = \eta(t)\Psi(t)$$

⇒ Time-dependent Dyson relation

$$h(t) = \eta(t)H(t)\eta^{-1}(t) + i\hbar\partial_t\eta(t)\eta^{-1}(t)$$

Theoretical framework (key equations):

Time-dependent Schrödinger eqn for $h(t) = h^\dagger(t)$, $H(t) \neq H^\dagger(t)$

$$h(t)\phi(t) = i\hbar\partial_t\phi(t), \quad \text{and} \quad H(t)\Psi(t) = i\hbar\partial_t\Psi(t)$$

Time-dependent Dyson operator

$$\phi(t) = \eta(t)\Psi(t)$$

⇒ Time-dependent Dyson relation

$$h(t) = \eta(t)H(t)\eta^{-1}(t) + i\hbar\partial_t\eta(t)\eta^{-1}(t)$$

⇒ Time-dependent quasi-Hermiticity relation

$$H^\dagger\rho(t) - \rho(t)H = i\hbar\partial_t\rho(t)$$

[from conjugating Dyson relation and $\rho(t) := \eta^\dagger(t)\eta(t)$]

Theoretical framework (interpretation):

Observables $o(t)$ in the Hermitian system are self-adjoint.

Observables $\mathcal{O}(t)$ in the non-Hermitian $\mathcal{O}(t)$ are quasi Hermitian

$$o(t) = \eta(t)\mathcal{O}(t)\eta^{-1}(t).$$

Theoretical framework (interpretation):

Observables $o(t)$ in the Hermitian system are self-adjoint.

Observables $\mathcal{O}(t)$ in the non-Hermitian $\mathcal{O}(t)$ are quasi Hermitian

$$o(t) = \eta(t)\mathcal{O}(t)\eta^{-1}(t).$$

Then we have

$$\langle \phi(t) | o(t) \phi(t) \rangle = \langle \Psi(t) | \rho(t) \mathcal{O}(t) \Psi(t) \rangle .$$

Theoretical framework (interpretation):

Observables $o(t)$ in the Hermitian system are self-adjoint.

Observables $\mathcal{O}(t)$ in the non-Hermitian $\mathcal{O}(t)$ are quasi Hermitian

$$o(t) = \eta(t)\mathcal{O}(t)\eta^{-1}(t).$$

Then we have

$$\langle \phi(t) | o(t) \phi(t) \rangle = \langle \Psi(t) | \rho(t) \mathcal{O}(t) \Psi(t) \rangle.$$

Since $H(t)$ is not quasi/pseudo Hermitian it is not an observable.

Instead the observable energy operator is

$$\tilde{H}(t) = \eta^{-1}(t)h(t)\eta(t) = H(t) + i\hbar\eta^{-1}(t)\partial_t\eta(t).$$

$H(t)$ is simply the Hamiltonian satisfying the TDSE and governing the evolution in time.

Unitary time-evolution:

Hermitian:

$$\phi(t) = u(t, t')\phi(t'), \quad u(t, t') = T \exp \left[-i \int_{t'}^t ds h(s) \right]$$

with

$$h(t)u(t, t') = i\hbar\partial_t u(t, t'), \quad u(t, t')u(t', t'') = u(t, t''), \quad u(t, t) = \mathbb{I}$$

$$\langle u(t, t')\phi(t') | u(t, t')\tilde{\phi}(t') \rangle = \langle \phi(t) | \tilde{\phi}(t) \rangle$$

Relation between $u(t, t')$ and $U(t, t')$:

$$U(t, t') = \eta^{-1}(t)u(t, t')\eta(t')$$

or the generalized Duhamel's formula

$$\begin{aligned} U(t, t') &= u(t, t') - \int_{t'}^t \frac{d}{ds} [U(t, s)u(s, t')] ds \\ &= u(t, t') - i\hbar \int_{t'}^t U(t, s) [H(s) - h(s)] u(s, t') ds \end{aligned}$$

Relation between $u(t, t')$ and $U(t, t')$:

$$U(t, t') = \eta^{-1}(t)u(t, t')\eta(t')$$

or the generalized Duhamel's formula

$$\begin{aligned} U(t, t') &= u(t, t') - \int_{t'}^t \frac{d}{ds} [U(t, s)u(s, t')] ds \\ &= u(t, t') - i\hbar \int_{t'}^t U(t, s) [H(s) - h(s)] u(s, t') ds \end{aligned}$$

Relation between Green's functions:

$$G_h(t, t') := -iu(t, t')\theta(t - t') \quad G_H(t, t') := -iU(t, t')\theta(t - t')$$

$$G_U(t, t') = G_u(t, t') + i \int_{-\infty}^{\infty} G_U(t, s) [H(s) - h(s)] G_u(s, t') ds$$

Ising quantum spin chain of length N

$$\mathcal{H} = -\frac{1}{2} \sum_{i=1}^N (\sigma_i^z + \lambda \sigma_i^x \sigma_{i+1}^x + i\kappa \sigma_i^x) \quad \kappa, \lambda \in \mathbb{R}$$

in a magnetic field in the z-direction and in a longitudinal imaginary field in the x-direction

- \mathcal{H} acts on the Hilbert space of the form $(\mathbb{C}^2)^{\otimes N}$
- $\sigma_i^{x,y,z} := \mathbb{I} \otimes \mathbb{I} \otimes \dots \otimes \sigma_i^{x,y,z} \otimes \dots \otimes \mathbb{I} \otimes \mathbb{I}$

$$\sigma^x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

Ising quantum spin chain of length N

$$\mathcal{H} = -\frac{1}{2} \sum_{i=1}^N (\sigma_i^z + \lambda \sigma_i^x \sigma_{i+1}^x + i\kappa \sigma_i^x) \quad \kappa, \lambda \in \mathbb{R}$$

in a magnetic field in the z-direction and in a longitudinal imaginary field in the x-direction

- \mathcal{H} acts on the Hilbert space of the form $(\mathbb{C}^2)^{\otimes N}$
- $\sigma_i^{x,y,z} := \mathbb{I} \otimes \mathbb{I} \otimes \dots \otimes \sigma_i^{x,y,z} \otimes \dots \otimes \mathbb{I} \otimes \mathbb{I}$

$$\sigma^x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

- \mathcal{H} is a perturbation of the $\mathcal{M}_{5,2}$ -model ($c=-22/5$)
in the $\mathcal{M}_{p,q}$ -series of minimal conformal field theories

Ising quantum spin chain of length N

$$\mathcal{H} = -\frac{1}{2} \sum_{i=1}^N (\sigma_i^z + \lambda \sigma_i^x \sigma_{i+1}^x + i\kappa \sigma_i^x) \quad \kappa, \lambda \in \mathbb{R}$$

in a magnetic field in the z-direction and in a longitudinal imaginary field in the x-direction

- \mathcal{H} acts on the Hilbert space of the form $(\mathbb{C}^2)^{\otimes N}$
- $\sigma_i^{x,y,z} := \mathbb{I} \otimes \mathbb{I} \otimes \dots \otimes \sigma_i^{x,y,z} \otimes \dots \otimes \mathbb{I} \otimes \mathbb{I}$

$$\sigma^x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

- \mathcal{H} is a perturbation of the $\mathcal{M}_{5,2}$ -model ($c=-22/5$)
in the $\mathcal{M}_{p,q}$ -series of minimal conformal field theories
- non-unitary for $p - q > 1 \Rightarrow$ non-Hermitian Hamiltonians

Ising quantum spin chain of length N

$$\mathcal{H} = -\frac{1}{2} \sum_{i=1}^N (\sigma_i^z + \lambda \sigma_i^x \sigma_{i+1}^x + i\kappa \sigma_i^x) \quad \kappa, \lambda \in \mathbb{R}$$

in a magnetic field in the z-direction and in a longitudinal imaginary field in the x-direction

- \mathcal{H} acts on the Hilbert space of the form $(\mathbb{C}^2)^{\otimes N}$
- $\sigma_i^{x,y,z} := \mathbb{I} \otimes \mathbb{I} \otimes \dots \otimes \sigma_i^{x,y,z} \otimes \dots \otimes \mathbb{I} \otimes \mathbb{I}$

$$\sigma^x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

- \mathcal{H} is a perturbation of the $\mathcal{M}_{5,2}$ -model ($c=-22/5$)
in the $\mathcal{M}_{p,q}$ -series of minimal conformal field theories
- non-unitary for $p - q > 1 \Rightarrow$ non-Hermitian Hamiltonians

[G. von Gehlen, J. Phys. A24 (1991) 5371]



\mathcal{PT} -symmetry for spin chains

\mathcal{PT} -symmetry for spin chains

- "macro-reflections": [Korff, Weston, J. Phys. A40 (2007)]

$$\mathcal{P}' : \sigma_i^{x,y,z} \rightarrow \sigma_{N+1-i}^{x,y,z}$$

$$\mathcal{P}' : \begin{array}{cccccccc} \nearrow_1 & - & - & \searrow_2 & - & - & \nwarrow_3 & - & \dots & - & - & \nearrow_{N-2} & - & - & \nearrow_{N-1} & - & - & \swarrow_N \end{array}$$

\mathcal{PT} -symmetry for spin chains

- "macro-reflections": [Korff, Weston, J. Phys. A40 (2007)]

$$\mathcal{P}' : \sigma_i^{x,y,z} \rightarrow \sigma_{N+1-i}^{x,y,z}$$



\mathcal{PT} -symmetry for spin chains

- "macro-reflections": [Korff, Weston, J. Phys. A40 (2007)]

$$\mathcal{P}' : \sigma_i^{x,y,z} \rightarrow \sigma_{N+1-i}^{x,y,z}$$



- but with $\mathcal{T} : i \rightarrow -i$ $[\mathcal{P}'\mathcal{T}, \mathcal{H}] \neq 0$

\mathcal{PT} -symmetry for spin chains

- "macro-reflections": [Korff, Weston, J. Phys. A40 (2007)]

$$\mathcal{P}' : \sigma_i^{x,y,z} \rightarrow \sigma_{N+1-i}^{x,y,z}$$



- but with $\mathcal{T} : i \rightarrow -i$ $[\mathcal{P}'\mathcal{T}, \mathcal{H}] \neq 0$

- "site-by-site reflections":

[Castro-Alvaredo, A.F., J.Phys. A42 (2009) 465211]

$$\mathcal{P} = \prod_{i=1}^N \sigma_i^z, \quad \text{with} \quad \mathcal{P}^2 = \mathbb{I}^{\otimes N}$$

$$\mathcal{P} : (\sigma_i^x, \sigma_i^y, \sigma_i^z) \rightarrow (-\sigma_i^x, -\sigma_i^y, \sigma_i^z)$$



\mathcal{PT} -symmetry for spin chains

- "macro-reflections": [Korff, Weston, J. Phys. A40 (2007)]

$$\mathcal{P}' : \sigma_i^{x,y,z} \rightarrow \sigma_{N+1-i}^{x,y,z}$$



- but with $\mathcal{T} : i \rightarrow -i$ $[\mathcal{P}'\mathcal{T}, \mathcal{H}] \neq 0$

- "site-by-site reflections":

[Castro-Alvaredo, A.F., J.Phys. A42 (2009) 465211]

$$\mathcal{P} = \prod_{i=1}^N \sigma_i^z, \quad \text{with} \quad \mathcal{P}^2 = \mathbb{I}^{\otimes N}$$

$$\mathcal{P} : (\sigma_i^x, \sigma_i^y, \sigma_i^z) \rightarrow (-\sigma_i^x, -\sigma_i^y, \sigma_i^z)$$



\mathcal{PT} -symmetry for spin chains

- "macro-reflections": [Korff, Weston, J. Phys. A40 (2007)]

$$\mathcal{P}' : \sigma_i^{x,y,z} \rightarrow \sigma_{N+1-i}^{x,y,z}$$



- but with $\mathcal{T} : i \rightarrow -i$ $[\mathcal{P}'\mathcal{T}, \mathcal{H}] \neq 0$

- "site-by-site reflections":

[Castro-Alvaredo, A.F., J.Phys. A42 (2009) 465211]

$$\mathcal{P} = \prod_{i=1}^N \sigma_i^z, \quad \text{with} \quad \mathcal{P}^2 = \mathbb{I}^{\otimes N}$$

$$\mathcal{P} : (\sigma_i^x, \sigma_i^y, \sigma_i^z) \rightarrow (-\sigma_i^x, -\sigma_i^y, \sigma_i^z)$$



$$\Rightarrow [\mathcal{PT}, \mathcal{H}] = 0$$

- Alternative definitions for parity:

$$\mathcal{P}_x := \prod_{i=1}^N \sigma_i^x \qquad \mathcal{P}_y := \prod_{i=1}^N \sigma_i^y$$

$$\mathcal{P}_x : (\sigma_i^x, \sigma_i^y, \sigma_i^z) \rightarrow (\sigma_i^x, -\sigma_i^y, -\sigma_i^z)$$

$$\mathcal{P}_y : (\sigma_i^x, \sigma_i^y, \sigma_i^z) \rightarrow (-\sigma_i^x, \sigma_i^y, -\sigma_i^z)$$

$$[\mathcal{PT}, \mathcal{H}] = 0, \quad [\mathcal{P}_x \mathcal{T}, \mathcal{H}] \neq 0, \quad [\mathcal{P}_y \mathcal{T}, \mathcal{H}] \neq 0, \quad [\mathcal{P}' \mathcal{T}, \mathcal{H}] \neq 0$$

- Alternative definitions for parity:

$$\mathcal{P}_x := \prod_{i=1}^N \sigma_i^x \quad \mathcal{P}_y := \prod_{i=1}^N \sigma_i^y$$

$$\mathcal{P}_x : (\sigma_i^x, \sigma_i^y, \sigma_i^z) \rightarrow (\sigma_i^x, -\sigma_i^y, -\sigma_i^z)$$

$$\mathcal{P}_y : (\sigma_i^x, \sigma_i^y, \sigma_i^z) \rightarrow (-\sigma_i^x, \sigma_i^y, -\sigma_i^z)$$

$$[\mathcal{PT}, \mathcal{H}] = 0, [\mathcal{P}_x \mathcal{T}, \mathcal{H}] \neq 0, [\mathcal{P}_y \mathcal{T}, \mathcal{H}] \neq 0, [\mathcal{P}' \mathcal{T}, \mathcal{H}] \neq 0$$

- XXZ-spin-chain in a magnetic field

$$\mathcal{H}_{\text{XXZ}} = \frac{1}{2} \sum_{i=1}^{N-1} [(\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y + \Delta_+ (\sigma_i^z \sigma_{i+1}^z - 1))] + \frac{\Delta_-}{2} (\sigma_1^z - \sigma_N^z),$$

$$\Delta_{\pm} = (q \pm q^{-1})/2 \quad \Rightarrow \mathcal{H}_{\text{XXZ}}^{\dagger} \neq \mathcal{H}_{\text{XXZ}} \text{ for } q \notin \mathbb{R}$$

$$[\mathcal{PT}, \mathcal{H}_{\text{XXZ}}] \neq 0 \quad [\mathcal{P}_x \mathcal{T}, \mathcal{H}_{\text{XXZ}}] = 0 \quad [\mathcal{P}_y \mathcal{T}, \mathcal{H}_{\text{XXZ}}] = 0 \quad [\mathcal{P}' \mathcal{T}, \mathcal{H}_{\text{XXZ}}] = 0$$

- Alternative definitions for parity:

$$\mathcal{P}_x := \prod_{i=1}^N \sigma_i^x \quad \mathcal{P}_y := \prod_{i=1}^N \sigma_i^y$$

$$\mathcal{P}_x : (\sigma_i^x, \sigma_i^y, \sigma_i^z) \rightarrow (\sigma_i^x, -\sigma_i^y, -\sigma_i^z)$$

$$\mathcal{P}_y : (\sigma_i^x, \sigma_i^y, \sigma_i^z) \rightarrow (-\sigma_i^x, \sigma_i^y, -\sigma_i^z)$$

$$[\mathcal{PT}, \mathcal{H}] = 0, [\mathcal{P}_x \mathcal{T}, \mathcal{H}] \neq 0, [\mathcal{P}_y \mathcal{T}, \mathcal{H}] \neq 0, [\mathcal{P}' \mathcal{T}, \mathcal{H}] \neq 0$$

- XXZ-spin-chain in a magnetic field

$$\mathcal{H}_{\text{XXZ}} = \frac{1}{2} \sum_{i=1}^{N-1} [(\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y + \Delta_+ (\sigma_i^z \sigma_{i+1}^z - 1))] + \frac{\Delta_-}{2} (\sigma_1^z - \sigma_N^z),$$

$$\Delta_{\pm} = (q \pm q^{-1})/2 \quad \Rightarrow \mathcal{H}_{\text{XXZ}}^{\dagger} \neq \mathcal{H}_{\text{XXZ}} \text{ for } q \notin \mathbb{R}$$

$$[\mathcal{PT}, \mathcal{H}_{\text{XXZ}}] \neq 0 \quad [\mathcal{P}_x \mathcal{T}, \mathcal{H}_{\text{XXZ}}] = 0 \quad [\mathcal{P}_y \mathcal{T}, \mathcal{H}_{\text{XXZ}}] = 0 \quad [\mathcal{P}' \mathcal{T}, \mathcal{H}_{\text{XXZ}}] = 0$$

These possibilities reflect the ambiguities in the observables.

\mathcal{PT} -symmetry \Rightarrow domains in the parameter space of λ and κ

Broken and unbroken \mathcal{PT} -symmetry

$$[\mathcal{PT}, \mathcal{H}] = 0 \quad \bigwedge \quad \mathcal{PT}\Phi(\lambda, \kappa) \begin{cases} = \Phi(\lambda, \kappa) & \text{for } (\lambda, \kappa) \in U_{\mathcal{PT}} \\ \neq \Phi(\lambda, \kappa) & \text{for } (\lambda, \kappa) \in U_{b\mathcal{PT}} \end{cases}$$

$(\lambda, \kappa) \in U_{\mathcal{PT}} \Rightarrow$ real eigenvalues

$(\lambda, \kappa) \in U_{b\mathcal{PT}} \Rightarrow$ eigenvalues in complex conjugate pairs

- The two site Hamiltonian

$$\begin{aligned}
 \mathcal{H} &= -\frac{1}{2} [\sigma_1^z + \sigma_2^z + 2\lambda\sigma_1^x\sigma_2^x + i\kappa(\sigma_2^x + \sigma_1^x)] \\
 &= -\frac{1}{2} [\sigma^z \otimes \mathbb{I} + \mathbb{I} \otimes \sigma^z + 2\lambda\sigma^x \otimes \sigma^x + i\kappa(\mathbb{I} \otimes \sigma^x + \sigma^x \otimes \mathbb{I})] \\
 &= - \begin{pmatrix} -1 & \frac{i\kappa}{2} & \frac{i\kappa}{2} & \lambda \\ \frac{i\kappa}{2} & 0 & \lambda & \frac{i\kappa}{2} \\ \frac{i\kappa}{2} & \lambda & 0 & \frac{i\kappa}{2} \\ \lambda & \frac{i\kappa}{2} & \frac{i\kappa}{2} & -1 \end{pmatrix}
 \end{aligned}$$

- The two site Hamiltonian

$$\begin{aligned}
 \mathcal{H} &= -\frac{1}{2} [\sigma_1^z + \sigma_2^z + 2\lambda\sigma_1^x\sigma_2^x + i\kappa(\sigma_2^x + \sigma_1^x)] \\
 &= -\frac{1}{2} [\sigma^z \otimes \mathbb{I} + \mathbb{I} \otimes \sigma^z + 2\lambda\sigma^x \otimes \sigma^x + i\kappa(\mathbb{I} \otimes \sigma^x + \sigma^x \otimes \mathbb{I})] \\
 &= - \begin{pmatrix} -1 & \frac{i\kappa}{2} & \frac{i\kappa}{2} & \lambda \\ \frac{i\kappa}{2} & 0 & \lambda & \frac{i\kappa}{2} \\ \frac{i\kappa}{2} & \lambda & 0 & \frac{i\kappa}{2} \\ \lambda & \frac{i\kappa}{2} & \frac{i\kappa}{2} & -1 \end{pmatrix}
 \end{aligned}$$

with periodic boundary condition $\sigma_{N+1}^x = \sigma_1^x$

- The two site Hamiltonian

$$\begin{aligned}
 \mathcal{H} &= -\frac{1}{2} [\sigma_1^z + \sigma_2^z + 2\lambda\sigma_1^x\sigma_2^x + i\kappa(\sigma_2^x + \sigma_1^x)] \\
 &= -\frac{1}{2} [\sigma^z \otimes \mathbb{I} + \mathbb{I} \otimes \sigma^z + 2\lambda\sigma^x \otimes \sigma^x + i\kappa(\mathbb{I} \otimes \sigma^x + \sigma^x \otimes \mathbb{I})] \\
 &= - \begin{pmatrix} -1 & \frac{i\kappa}{2} & \frac{i\kappa}{2} & \lambda \\ \frac{i\kappa}{2} & 0 & \lambda & \frac{i\kappa}{2} \\ \frac{i\kappa}{2} & \lambda & 0 & \frac{i\kappa}{2} \\ \lambda & \frac{i\kappa}{2} & \frac{i\kappa}{2} & -1 \end{pmatrix}
 \end{aligned}$$

with periodic boundary condition $\sigma_{N+1}^x = \sigma_1^x$

- domain of unbroken \mathcal{PT} -symmetry:

- The two site Hamiltonian

$$\begin{aligned}
 \mathcal{H} &= -\frac{1}{2} [\sigma_1^z + \sigma_2^z + 2\lambda\sigma_1^x\sigma_2^x + i\kappa(\sigma_2^x + \sigma_1^x)] \\
 &= -\frac{1}{2} [\sigma^z \otimes \mathbb{I} + \mathbb{I} \otimes \sigma^z + 2\lambda\sigma^x \otimes \sigma^x + i\kappa(\mathbb{I} \otimes \sigma^x + \sigma^x \otimes \mathbb{I})] \\
 &= - \begin{pmatrix} -1 & \frac{i\kappa}{2} & \frac{i\kappa}{2} & \lambda \\ \frac{i\kappa}{2} & 0 & \lambda & \frac{i\kappa}{2} \\ \frac{i\kappa}{2} & \lambda & 0 & \frac{i\kappa}{2} \\ \lambda & \frac{i\kappa}{2} & \frac{i\kappa}{2} & -1 \end{pmatrix}
 \end{aligned}$$

with periodic boundary condition $\sigma_{N+1}^x = \sigma_1^x$

- domain of unbroken \mathcal{PT} -symmetry:


char. polynomial factorises into 1st and 3rd order

discriminant: $\Delta = r^2 - q^3$

$$q = \frac{1}{9} (-3\kappa^2 + 4\lambda^2 + 3), \quad r = \frac{\lambda}{27} (18\kappa^2 + 8\lambda^2 + 9)$$

Deformed quantum spin chains (Exact Results, $N = 2$)

$$U_{PT} = \left\{ \lambda, \kappa : \kappa^6 + 8\lambda^2\kappa^4 - 3\kappa^4 + 16\lambda^4\kappa^2 + 20\lambda^2\kappa^2 + 3\kappa^2 - \lambda^2 \leq 1 \right\}$$





- Right eigenvectors of \mathcal{H} :

$$|\Phi_1\rangle = (0, -1, -1, 0) \quad |\Phi_n\rangle = (\gamma_n, -\alpha_n, -\alpha_n, \beta_n) \quad n = 2, 3, 4$$

$$\alpha_n = i\kappa(\lambda - \varepsilon_n + 1)$$

$$\beta_n = \kappa^2 + 2\lambda^2 + 2\lambda\varepsilon_n$$

$$\gamma_n = -\kappa^2 - 2\varepsilon_n^2 + 2\lambda - 2\lambda\varepsilon_n + 2\varepsilon_n$$

- Right eigenvectors of \mathcal{H} :

$$|\Phi_1\rangle = (0, -1, -1, 0) \quad |\Phi_n\rangle = (\gamma_n, -\alpha_n, -\alpha_n, \beta_n) \quad n = 2, 3, 4$$

$$\alpha_n = i\kappa(\lambda - \varepsilon_n + 1)$$

$$\beta_n = \kappa^2 + 2\lambda^2 + 2\lambda\varepsilon_n$$

$$\gamma_n = -\kappa^2 - 2\varepsilon_n^2 + 2\lambda - 2\lambda\varepsilon_n + 2\varepsilon_n$$

- signature: $\mathbf{s} = (+, -, +, -)$

$$\mathcal{P} |\Phi_n\rangle = \mathbf{s}_n |\Psi_n\rangle$$

from relating left and right eigenvectors

- \mathcal{C} -operator:

$$\mathcal{C} = \sum_n s_n |\Phi_n\rangle \langle \Psi_n|$$

$$= \begin{pmatrix} C_5 & -C_3 & -C_3 & C_4 \\ -C_3 & -C_1 - 1 & -C_1 & C_2 \\ -C_3 & -C_1 & -C_1 - 1 & C_2 \\ C_4 & C_2 & C_2 & 2(C_1 + 1) - C_5 \end{pmatrix}$$

$$C_1 = \frac{\alpha_4^2}{N_4^2} - \frac{\alpha_2^2}{N_2^2} - \frac{\alpha_3^2}{N_3^2} - \frac{1}{2}, \quad C_2 = \frac{\alpha_4\beta_4}{N_4^2} - \frac{\alpha_2\beta_2}{N_2^2} - \frac{\alpha_3\beta_3}{N_3^2},$$

$$C_3 = \frac{\alpha_2\gamma_2}{N_2^2} + \frac{\alpha_3\gamma_3}{N_3^2} - \frac{\alpha_4\gamma_4}{N_4^2}, \quad C_4 = \frac{\beta_2\gamma_2}{N_2^2} + \frac{\beta_3\gamma_3}{N_3^2} - \frac{\beta_4\gamma_4}{N_4^2},$$

$$C_5 = \frac{\gamma_2^2}{N_2^2} + \frac{\gamma_3^2}{N_3^2} - \frac{\gamma_4^2}{N_4^2}$$

$$N_1 = \sqrt{2}, \quad N_n = \sqrt{2\alpha_n^2 + \beta_n^2 + \gamma_n^2} \quad \text{for } n = 2, 3, 4$$

- metric operator:

$$\rho = \mathcal{P}\mathcal{C} = \begin{pmatrix} C_5 & -C_3 & -C_3 & C_4 \\ C_3 & 1 + C_1 & C_1 & -C_2 \\ C_3 & C_1 & 1 + C_1 & -C_2 \\ C_4 & C_2 & C_2 & 2(1 + C_1) - C_5 \end{pmatrix}$$

- since $i\alpha_i, \beta_i, \gamma_i \in \mathbb{R}$
 $\Rightarrow C_1, iC_2, iC_3, C_4, C_5 \in \mathbb{R}$
 $\Rightarrow \rho$ is Hermitian $\rho = \rho^\dagger$

- metric operator:

$$\rho = \mathcal{P}\mathcal{C} = \begin{pmatrix} C_5 & -C_3 & -C_3 & C_4 \\ C_3 & 1 + C_1 & C_1 & -C_2 \\ C_3 & C_1 & 1 + C_1 & -C_2 \\ C_4 & C_2 & C_2 & 2(1 + C_1) - C_5 \end{pmatrix}$$

- since $i\alpha_i, \beta_i, \gamma_i \in \mathbb{R}$
 $\Rightarrow C_1, iC_2, iC_3, C_4, C_5 \in \mathbb{R}$
 $\Rightarrow \rho$ is Hermitian $\rho = \rho^\dagger$
- EV of ρ :

$$y_1 = y_2 = 1, \quad y_{3/4} = 1 + 2C_1 \pm 2\sqrt{C_1(1 + C_1)}$$

- metric operator:

$$\rho = \mathcal{P}\mathcal{C} = \begin{pmatrix} C_5 & -C_3 & -C_3 & C_4 \\ C_3 & 1 + C_1 & C_1 & -C_2 \\ C_3 & C_1 & 1 + C_1 & -C_2 \\ C_4 & C_2 & C_2 & 2(1 + C_1) - C_5 \end{pmatrix}$$

- since $i\alpha_i, \beta_i, \gamma_i \in \mathbb{R}$
 $\Rightarrow C_1, iC_2, iC_3, C_4, C_5 \in \mathbb{R}$
 $\Rightarrow \rho$ is Hermitian $\rho = \rho^\dagger$
- EV of ρ :

$$y_1 = y_2 = 1, \quad y_{3/4} = 1 + 2C_1 \pm 2\sqrt{C_1(1 + C_1)}$$

since $C_1 > 0 \Rightarrow \rho$ is positive

- square root of the metric operator:

$$\eta = \rho^{1/2} = UD^{1/2}U^{-1}$$

where $D = \text{diag}(y_1, y_2, y_3, y_4)$, $U = \{r_1, r_2, r_3, r_4\}$

$$|r_1\rangle = (0, -1, 1, 0)$$

$$|r_2\rangle = (C_4, 0, 0, 1 - C_5),$$

$$|r_{3/4}\rangle = (\tilde{\gamma}_{3/4}, \tilde{\alpha}_{3/4}, \tilde{\alpha}_{3/4}, \tilde{\beta}_{3/4})$$

$$\tilde{\alpha}_{3/4} = y_{3/4}(C_3 C_4 + C_2(-4C_1 + C_5 - 1))/2 - C_3 C_4$$

$$\tilde{\beta}_{3/4} = -C_3^2 - C_1 - C_1 C_5 + (C_3^2 + C_1(4C_1 - C_5 + 3)) y_{3/4},$$

$$\tilde{\gamma}_{3/4} = C_1 C_4 - C_2 C_3 + (C_2 C_3 + C_1 C_4) y_{3/4}$$


- isospectral Hermitian counterpart:

$$\begin{aligned}
 h &= \eta \mathcal{H} \eta^{-1} \\
 &= \mu_1 \sigma_x \otimes \sigma_x + \mu_2 \sigma_y \otimes \sigma_y + \mu_3 \sigma_z \otimes \sigma_z + \mu_4 (\sigma_z \otimes \mathbb{I} + \mathbb{I} \otimes \sigma_z)
 \end{aligned}$$

$$\mu_1, \mu_2, \mu_3, \mu_4 \in \mathbb{R}$$

for $\lambda = 0.1, \kappa = 0.5$:

$$h = \begin{pmatrix} -0.829536 & 0 & 0 & -0.0606492 \\ 0 & -0.0341687 & -0.1341687 & 0 \\ 0 & -0.1341687 & -0.0341687 & 0 \\ -0.0606492 & 0 & 0 & 0.897873 \end{pmatrix}$$

The magnetization in the z-direction for $N = 2$:

Deformed quantum spin chains ($N \neq 2$, perturbation theory)

- Perturbation theory about the Hermitian part

$$H(\lambda, \kappa) = h_0(\lambda) + i\kappa h_1 \quad h_0 = h_0^\dagger, h_1 = h_1^\dagger \quad \kappa \in \mathbb{R}$$

assume $\eta = \eta^\dagger = e^{q/2} \Rightarrow$ solve for q

$$H^\dagger = e^q H e^{-q} = H + [q, H] + \frac{1}{2}[q, [q, H]] + \frac{1}{3!}[q, [q, [q, H]]] + \dots$$

for $c_q^{(\ell+1)}(h_0) = [q, \dots [q, [q, h_0]] \dots] = 0$ closed formulae:

$$h = h_0 + \sum_{n=1}^{\lfloor \frac{\ell}{2} \rfloor} \frac{(-1)^n E_n}{4^n (2n)!} c_q^{(2n)}(h_0) \quad H = h_0 - \sum_{n=1}^{\lfloor \frac{\ell+1}{2} \rfloor} \frac{\kappa_{2n-1}}{(2n-1)!} c_q^{(2n-1)}(h_0)$$

$E_n \equiv$ Euler numbers, e.g. $E_1 = 1, E_2 = 5, E_3 = 61, \dots$

$$\kappa_n = \frac{1}{2^n} \sum_{m=1}^{\lfloor (n+1)/2 \rfloor} (-1)^{n+m} \binom{n}{2m} E_m$$

$\kappa_1 = 1/2, \kappa_3 = -1/4, \kappa_5 = 1/2, \kappa_7 = -17/8, \dots$

[C. F. de Morisson Faria, A.F., J. Phys. A39 (2006) 9269]

exact result for $N = 2$:

$\lambda = 0.1, \kappa = 0.5$:

$$h = \begin{pmatrix} -0.829536 & 0 & 0 & -0.0606492 \\ 0 & -0.0341687 & -0.1341687 & 0 \\ 0 & -0.1341687 & -0.0341687 & 0 \\ -0.0606492 & 0 & 0 & 0.897873 \end{pmatrix}$$

$\lambda = 0.9, \kappa = 0.1$:

$$h = \begin{pmatrix} -0.985439 & 0 & 0 & -0.890532 \\ 0 & -0.0094167 & -0.909417 & 0 \\ 0 & -0.909417 & -0.0094167 & 0 \\ -0.890532 & 0 & 0 & 1.00427 \end{pmatrix}$$

perturbative result 4th order for $N = 2$:

$\lambda = 0.1, \kappa = 0.5$:

$$h = \begin{pmatrix} -0.829534 & 0 & 0 & -0.0606716 \\ 0 & -0.0341688 & -0.134169 & 0 \\ 0 & -0.134169 & -0.0341688 & 0 \\ -0.0606716 & 0 & 0 & 0.897872 \end{pmatrix}$$

$\lambda = 0.9, \kappa = 0.1$:

$$h = \begin{pmatrix} -0.985439 & 0 & 0 & -0.890532 \\ 0 & -0.0094167 & -0.909417 & 0 \\ 0 & -0.909417 & -0.0094167 & 0 \\ -0.890532 & 0 & 0 & 1.00427 \end{pmatrix}$$

Deformed quantum spin chains ($N \neq 2$, perturbation theory)

- new notation:

$$S_{a_1 a_2 \dots a_p}^N := \sum_{k=1}^N \sigma_k^{a_1} \sigma_{k+1}^{a_2} \dots \sigma_{k+p-1}^{a_p}, \quad a_i = x, y, z, u; i = 1, \dots, p \leq N$$

with $\sigma^u = \mathbb{I}$ to allow for non-local interactions

- for instance:

$$\begin{aligned} H(\lambda, \kappa) &= -\frac{1}{2} \sum_{j=1}^N (\sigma_j^z + \lambda \sigma_j^x \sigma_{j+1}^x + i\kappa \sigma_j^x), \quad \lambda, \kappa \in \mathbb{R} \\ &= -\frac{1}{2} (\mathbf{S}_z^N + \lambda \mathbf{S}_{xx}^N) - i\kappa \frac{1}{2} \mathbf{S}_x^N \end{aligned}$$

- perturbative result for $N = 3$:

$$\begin{aligned} h &= \mu_{xx}^3(\lambda, \kappa) \mathbf{S}_{xx}^3 + \mu_{yy}^3(\lambda, \kappa) \mathbf{S}_{yy}^3 + \mu_{zz}^3(\lambda, \kappa) \mathbf{S}_{zz}^3 + \mu_z^3(\lambda, \kappa) \mathbf{S}_z^3 \\ &\quad + \mu_{xxz}^3(\lambda, \kappa) \mathbf{S}_{xxz}^3 + \mu_{yyz}^3(\lambda, \kappa) \mathbf{S}_{yyz}^3 + \mu_{zzz}^3(\lambda, \kappa) \mathbf{S}_{zzz}^3 \end{aligned}$$

- perturbative result for $N = 4$:

$$\begin{aligned}
 h = & \mu_{xx}^4(\lambda, \kappa) \mathbf{S}_{xx}^4 + \nu_{xx}^4(\lambda, \kappa) \mathbf{S}_{xux}^4 + \mu_{yy}^4(\lambda, \kappa) \mathbf{S}_{yy}^4 + \nu_{yy}^4(\lambda, \kappa) \mathbf{S}_{yuy}^4 \\
 & + \mu_{zz}^4(\lambda, \kappa) \mathbf{S}_{zz}^4 + \nu_{zz}^4(\lambda, \kappa) \mathbf{S}_{zuz}^4 + \mu_z^4(\lambda, \kappa) \mathbf{S}_z^4 + \mu_{xzx}^4(\lambda, \kappa) \mathbf{S}_{xzx}^4 \\
 & + \mu_{xxz}^4(\lambda, \kappa) (\mathbf{S}_{xxz}^4 + \mathbf{S}_{zxx}^4) + \mu_{yyz}^4(\lambda, \kappa) (\mathbf{S}_{yyz}^4 + \mathbf{S}_{zyy}^4) \\
 & + \mu_{yzy}^4(\lambda, \kappa) \mathbf{S}_{yzy}^4 + \mu_{zzz}^4(\lambda, \kappa) \mathbf{S}_{zzz}^4 + \mu_{xxxx}^4(\lambda, \kappa) \mathbf{S}_{xxxx}^4 \\
 & + \mu_{yyyy}^4(\lambda, \kappa) \mathbf{S}_{yyyy}^4 + \mu_{zzzz}^4(\lambda, \kappa) \mathbf{S}_{zzzz}^4 + \mu_{xxyy}^4(\lambda, \kappa) \mathbf{S}_{xxyy}^4 \\
 & + \mu_{xyxy}^4(\lambda, \kappa) \mathbf{S}_{xyxy}^4 + \mu_{zzyy}^4(\lambda, \kappa) \mathbf{S}_{zzyy}^4 + \mu_{zyzy}^4(\lambda, \kappa) \mathbf{S}_{zyzy}^4 \\
 & + \mu_{xxzz}^4(\lambda, \kappa) \mathbf{S}_{xxzz}^4 + \mu_{xzxz}^4(\lambda, \kappa) \mathbf{S}_{xzxz}^4
 \end{aligned}$$

non-local terms

Calogero-Moser-Sutherland models (extended)

$$\mathcal{H}_{BK} = \frac{p^2}{2} + \frac{\omega^2}{2} \sum_i q_i^2 + \frac{g^2}{2} \sum_{i \neq k} \frac{1}{(q_i - q_k)^2} + i\tilde{g} \sum_{i \neq k} \frac{1}{(q_i - q_k)} p_i$$

with $g, \tilde{g} \in \mathbb{R}, q, p \in \mathbb{R}^{\ell+1}$

[B. Basu-Mallick, A. Kundu, Phys. Rev. B62 (2000) 9927]

Calogero-Moser-Sutherland models (extended)

$$\mathcal{H}_{BK} = \frac{p^2}{2} + \frac{\omega^2}{2} \sum_i q_i^2 + \frac{g^2}{2} \sum_{i \neq k} \frac{1}{(q_i - q_k)^2} + i\tilde{g} \sum_{i \neq k} \frac{1}{(q_i - q_k)} p_i$$

with $g, \tilde{g} \in \mathbb{R}, q, p \in \mathbb{R}^{\ell+1}$

[B. Basu-Mallick, A. Kundu, Phys. Rev. B62 (2000) 9927]

- 1 Representation independent formulation?
- 2 Other potentials apart from the rational one?
- 3 Other algebras apart from A_n, B_n or Coxeter groups?
- 4 Is it possible to include more coupling constants?
- 5 Are the extensions still integrable?

- Generalize Hamiltonian to:

$$\mathcal{H}_\mu = \frac{1}{2}p^2 + \frac{1}{2} \sum_{\alpha \in \Delta} g_\alpha^2 V(\alpha \cdot q) + i\mu \cdot p$$

· Now Δ is any root system

· $\mu = 1/2 \sum_{\alpha \in \Delta} \tilde{g}_\alpha f(\alpha \cdot q)\alpha$, $f(x) = 1/x$ $V(x) = f^2(x)$

[A. F., Mod. Phys. Lett. A21 (2006) 691, Acta P. 47 (2007) 44]

- Generalize Hamiltonian to:

$$\mathcal{H}_\mu = \frac{1}{2}p^2 + \frac{1}{2} \sum_{\alpha \in \Delta} g_\alpha^2 V(\alpha \cdot q) + i\mu \cdot p$$

· Now Δ is any root system

· $\mu = 1/2 \sum_{\alpha \in \Delta} \tilde{g}_\alpha f(\alpha \cdot q) \alpha$, $f(x) = 1/x$ $V(x) = f^2(x)$

[A. F., Mod. Phys. Lett. A21 (2006) 691, Acta P. 47 (2007) 44]

- Not so obvious that one can re-write

$$\mathcal{H}_\mu = \frac{1}{2}(p+i\mu)^2 + \frac{1}{2} \sum_{\alpha \in \Delta} \hat{g}_\alpha^2 V(\alpha \cdot q), \quad \hat{g}_\alpha^2 = \begin{cases} g_s^2 + \alpha_s^2 \tilde{g}_s^2 & \alpha \in \Delta_s \\ g_l^2 + \alpha_l^2 \tilde{g}_l^2 & \alpha \in \Delta_l \end{cases}$$

- Generalize Hamiltonian to:

$$\mathcal{H}_\mu = \frac{1}{2}p^2 + \frac{1}{2} \sum_{\alpha \in \Delta} g_\alpha^2 V(\alpha \cdot q) + i\mu \cdot p$$

· Now Δ is any root system

· $\mu = 1/2 \sum_{\alpha \in \Delta} \tilde{g}_\alpha f(\alpha \cdot q) \alpha$, $f(x) = 1/x$ $V(x) = f^2(x)$

[A. F., Mod. Phys. Lett. A21 (2006) 691, Acta P. 47 (2007) 44]

- Not so obvious that one can re-write

$$\mathcal{H}_\mu = \frac{1}{2}(p+i\mu)^2 + \frac{1}{2} \sum_{\alpha \in \Delta} \hat{g}_\alpha^2 V(\alpha \cdot q), \quad \hat{g}_\alpha^2 = \begin{cases} g_s^2 + \alpha_s^2 \tilde{g}_s^2 & \alpha \in \Delta_s \\ g_l^2 + \alpha_l^2 \tilde{g}_l^2 & \alpha \in \Delta_l \end{cases}$$

$$\Rightarrow \mathcal{H}_\mu = \eta^{-1} h_{\text{Cal}} \eta \quad \text{with} \quad \eta = e^{-q \cdot \mu}$$

- Generalize Hamiltonian to:

$$\mathcal{H}_\mu = \frac{1}{2}p^2 + \frac{1}{2} \sum_{\alpha \in \Delta} g_\alpha^2 V(\alpha \cdot q) + i\mu \cdot p$$

· Now Δ is any root system

· $\mu = 1/2 \sum_{\alpha \in \Delta} \tilde{g}_\alpha f(\alpha \cdot q) \alpha$, $f(x) = 1/x$ $V(x) = f^2(x)$

[A. F., Mod. Phys. Lett. A21 (2006) 691, Acta P. 47 (2007) 44]

- Not so obvious that one can re-write

$$\mathcal{H}_\mu = \frac{1}{2}(p+i\mu)^2 + \frac{1}{2} \sum_{\alpha \in \Delta} \hat{g}_\alpha^2 V(\alpha \cdot q), \quad \hat{g}_\alpha^2 = \begin{cases} g_s^2 + \alpha_s^2 \tilde{g}_s^2 & \alpha \in \Delta_s \\ g_l^2 + \alpha_l^2 \tilde{g}_l^2 & \alpha \in \Delta_l \end{cases}$$

$$\Rightarrow \mathcal{H}_\mu = \eta^{-1} h_{\text{Cal}} \eta \quad \text{with} \quad \eta = e^{-q \cdot \mu}$$

- integrability follows trivially $\dot{L} = [L, M]: L(p) \rightarrow L(p + i\mu)$

Extended Calogero-Moser-Sutherland models

- Generalize Hamiltonian to:

$$\mathcal{H}_\mu = \frac{1}{2}p^2 + \frac{1}{2} \sum_{\alpha \in \Delta} g_\alpha^2 V(\alpha \cdot q) + i\mu \cdot p$$

· Now Δ is any root system

· $\mu = 1/2 \sum_{\alpha \in \Delta} \tilde{g}_\alpha f(\alpha \cdot q) \alpha$, $f(x) = 1/x$ $V(x) = f^2(x)$

[A. F., Mod. Phys. Lett. A21 (2006) 691, Acta P. 47 (2007) 44]

- Not so obvious that one can re-write

$$\mathcal{H}_\mu = \frac{1}{2}(p+i\mu)^2 + \frac{1}{2} \sum_{\alpha \in \Delta} \hat{g}_\alpha^2 V(\alpha \cdot q), \quad \hat{g}_\alpha^2 = \begin{cases} g_s^2 + \alpha_s^2 \tilde{g}_s^2 & \alpha \in \Delta_s \\ g_l^2 + \alpha_l^2 \tilde{g}_l^2 & \alpha \in \Delta_l \end{cases}$$

$$\Rightarrow \mathcal{H}_\mu = \eta^{-1} h_{\text{Cal}} \eta \quad \text{with} \quad \eta = e^{-q \cdot \mu}$$

- integrability follows trivially $\dot{L} = [L, M]: L(p) \rightarrow L(p + i\mu)$

- computing backwards for any CMS-potential

$$\mathcal{H}_\mu = \frac{1}{2}p^2 + \frac{1}{2} \sum_{\alpha \in \Delta} \hat{g}_\alpha^2 V(\alpha \cdot q) + i\mu \cdot p - \frac{1}{2}\mu^2$$

- $\mu^2 = \alpha_s^2 \tilde{g}_s^2 \sum_{\alpha \in \Delta_s} V(\alpha \cdot q) + \alpha_l^2 \tilde{g}_l^2 \sum_{\alpha \in \Delta_l} V(\alpha \cdot q)$ only for V rational

- From real fields to complex particle systems

- i) No restrictions

e.g. Benjamin-Ono equation

$$u_t + uu_x + \lambda Hu_{xx} = 0 \quad (*)$$

$H \equiv$ Hilbert transform, i.e. $Hu(x) = \frac{P}{\pi} \int_{-\infty}^{\infty} \frac{u(x)}{z-x} dz$

Then

$$u(x, t) = \frac{\lambda}{2} \sum_{k=1}^{\ell} \left(\frac{i}{x - z_k} - \frac{i}{x - z_k^*} \right) \in \mathbb{R}$$

satisfies (*) iff z_k obeys the A_n -Calogero equ. of motion

$$\ddot{z}_k = \frac{\lambda^2}{2} \sum_{k \neq j} (z_j - z_k)^{-3}$$

[H. Chen, N. Pereira, Phys. Fluids 22 (1979) 187]

[talk by J. Feinberg, PHHQP workshop VI, 2007, London]

ii) restrict to submanifold

Theorem: [Airault, McKean, Moser, CPAM, (1977) 95]

Given a Hamiltonian $H(x_1, \dots, x_n, \dot{x}_1, \dots, \dot{x}_n)$ with flow

$$\dot{x}_i = \partial H / \partial \dot{x}_i \quad \text{and} \quad \ddot{x}_i = -\partial H / \partial x_i \quad i = 1, \dots, n$$

and conserved charges I_j in involution with H , i.e.

$\{I_j, H\} = 0$. Then the locus of $\text{grad } I = 0$ is invariant.

Example: Boussinesq equation

$$v_{tt} = a(v^2)_{xx} + bv_{xxxx} + v_{xx} \quad (**)$$

Then

$$v(x, t) = c \sum_{k=1}^{\ell} (x - z_k)^{-2}$$


satisfies (**) iff $b=1/12$, $c=-a/2$ and z_k obeys

$$\ddot{z}_k = 2 \sum_{j \neq k} (z_j - z_k)^{-3} \quad \Leftrightarrow \quad \ddot{z}_k = -\frac{\partial H}{\partial z_i}$$

$$\dot{z}_k = 1 - \sum_{j \neq k} (z_j - z_k)^{-2} \quad \Leftrightarrow \quad \text{grad}(I_3 - I_1) = 0$$



Constrained field equations \rightarrow complex Calogero models



[P. Assis and A.F., J. Phys. A42 (2009) 425206]

Consider

Antilinearly invariant deformed Calogero model

$$\mathcal{H}_{\text{PTCMS}} = \frac{p^2}{2} + \frac{m^2}{16} \sum_{\alpha \in \Delta_s} (\alpha \cdot \tilde{q})^2 + \frac{1}{2} \sum_{\alpha \in \Delta} g_\alpha V(\alpha \cdot \tilde{q}), \quad m, g_\alpha \in \mathbb{R}$$

Define deformed coordinates (A_2)

$$q_1 \rightarrow \tilde{q}_1 = q_1 \cosh \varepsilon + i\sqrt{3}(q_2 - q_3) \sinh \varepsilon$$

$$q_2 \rightarrow \tilde{q}_2 = q_2 \cosh \varepsilon + i\sqrt{3}(q_3 - q_1) \sinh \varepsilon$$

$$q_3 \rightarrow \tilde{q}_3 = q_3 \cosh \varepsilon + i\sqrt{3}(q_1 - q_2) \sinh \varepsilon$$

With standard 3D representation for the simple A_2 -roots

$\alpha_1 = \{1, -1, 0\}$, $\alpha_2 = \{0, 1, -1\}$, $q_{ij} := q_i - q_j$ compute

$$\alpha_1 \cdot \tilde{q} = q_{12} \cosh \varepsilon - \frac{i}{\sqrt{3}}(q_{13} + q_{23}) \sinh \varepsilon,$$

$$\alpha_2 \cdot \tilde{q} = q_{23} \cosh \varepsilon - \frac{i}{\sqrt{3}}(q_{21} + q_{31}) \sinh \varepsilon,$$

$$(\alpha_1 + \alpha_2) \cdot \tilde{q} = q_{13} \cosh \varepsilon + \frac{i}{\sqrt{3}}(q_{12} + q_{32}) \sinh \varepsilon.$$

Define deformed coordinates (A_2)

$$q_1 \rightarrow \tilde{q}_1 = q_1 \cosh \varepsilon + i\sqrt{3}(q_2 - q_3) \sinh \varepsilon$$

$$q_2 \rightarrow \tilde{q}_2 = q_2 \cosh \varepsilon + i\sqrt{3}(q_3 - q_1) \sinh \varepsilon$$

$$q_3 \rightarrow \tilde{q}_3 = q_3 \cosh \varepsilon + i\sqrt{3}(q_1 - q_2) \sinh \varepsilon$$

With standard 3D representation for the simple A_2 -roots

$\alpha_1 = \{1, -1, 0\}$, $\alpha_2 = \{0, 1, -1\}$, $q_{ij} := q_i - q_j$ compute

$$\alpha_1 \cdot \tilde{q} = q_{12} \cosh \varepsilon - \frac{\iota}{\sqrt{3}}(q_{13} + q_{23}) \sinh \varepsilon,$$

$$\alpha_2 \cdot \tilde{q} = q_{23} \cosh \varepsilon - \frac{\iota}{\sqrt{3}}(q_{21} + q_{31}) \sinh \varepsilon,$$

$$(\alpha_1 + \alpha_2) \cdot \tilde{q} = q_{13} \cosh \varepsilon + \frac{\iota}{\sqrt{3}}(q_{12} + q_{32}) \sinh \varepsilon.$$

Symmetries:

$$\mathcal{S}_1 : \quad q_1 \leftrightarrow q_2, q_3 \leftrightarrow q_3, \iota \rightarrow -\iota,$$

$$\mathcal{S}_2 : \quad q_2 \leftrightarrow q_3, q_1 \leftrightarrow q_1, \iota \rightarrow -\iota.$$

Note, this Hamiltonian also results from deforming the roots:

$$\alpha_1 \rightarrow \tilde{\alpha}_1 = \alpha_1 \cosh \varepsilon + i\sqrt{3} \sinh \varepsilon \lambda_2$$

$$\alpha_2 \rightarrow \tilde{\alpha}_2 = \alpha_2 \cosh \varepsilon - i\sqrt{3} \sinh \varepsilon \lambda_1$$

Thus

$$\begin{aligned} \mathcal{H}_{PTCMS} &= \frac{p^2}{2} + \frac{m^2}{16} \sum_{\tilde{\alpha} \in \tilde{\Delta}_s} (\tilde{\alpha} \cdot q)^2 + \frac{1}{2} \sum_{\tilde{\alpha} \in \tilde{\Delta}} g_{\tilde{\alpha}} V(\tilde{\alpha} \cdot q), \quad m, g_{\tilde{\alpha}} \in \mathbb{R} \\ &= \frac{p^2}{2} + \frac{m^2}{16} \sum_{\alpha \in \Delta_s} (\alpha \cdot \tilde{q})^2 + \frac{1}{2} \sum_{\alpha \in \Delta} g_{\alpha} V(\alpha \cdot \tilde{q}), \quad m, g_{\alpha} \in \mathbb{R} \end{aligned}$$

Note, this Hamiltonian also results from deforming the roots:

$$\alpha_1 \rightarrow \tilde{\alpha}_1 = \alpha_1 \cosh \varepsilon + i\sqrt{3} \sinh \varepsilon \lambda_2$$

$$\alpha_2 \rightarrow \tilde{\alpha}_2 = \alpha_2 \cosh \varepsilon - i\sqrt{3} \sinh \varepsilon \lambda_1$$

Thus

$$\mathcal{H}_{\mathcal{PTCMS}} = \frac{p^2}{2} + \frac{m^2}{16} \sum_{\tilde{\alpha} \in \tilde{\Delta}_s} (\tilde{\alpha} \cdot q)^2 + \frac{1}{2} \sum_{\tilde{\alpha} \in \tilde{\Delta}} g_{\tilde{\alpha}} V(\tilde{\alpha} \cdot q), \quad m, g_{\tilde{\alpha}} \in \mathbb{R}$$

$$= \frac{p^2}{2} + \frac{m^2}{16} \sum_{\alpha \in \Delta_s} (\alpha \cdot \tilde{q})^2 + \frac{1}{2} \sum_{\alpha \in \Delta} g_{\alpha} V(\alpha \cdot \tilde{q}), \quad m, g_{\alpha} \in \mathbb{R}$$

Symmetries:

$$\sigma_1^{\varepsilon} : \tilde{\alpha}_1 \leftrightarrow -\tilde{\alpha}_1, \tilde{\alpha}_2 \leftrightarrow \tilde{\alpha}_1 + \tilde{\alpha}_2 \quad \Leftrightarrow \quad q_1 \leftrightarrow q_2, q_3 \leftrightarrow q_3, \iota \rightarrow -\iota$$

$$\sigma_2^{\varepsilon} : \tilde{\alpha}_2 \leftrightarrow -\tilde{\alpha}_2, \tilde{\alpha}_1 \leftrightarrow \tilde{\alpha}_1 + \tilde{\alpha}_2 \quad \Leftrightarrow \quad q_2 \leftrightarrow q_3, q_1 \leftrightarrow q_1, \iota \rightarrow -\iota$$

Construction of antilinear deformations

- Involution $\in \mathcal{W} \equiv$ Coxeter group \Rightarrow deform in antilinear way
- Find a linear deformation map:

$$\delta : \Delta \rightarrow \tilde{\Delta}(\varepsilon) \quad \alpha \mapsto \tilde{\alpha} = \theta_\varepsilon \alpha$$

$$\alpha_j \in \Delta \subset \mathbb{R}^n, \quad \tilde{\alpha}_j(\varepsilon) \in \tilde{\Delta}(\varepsilon) \subset \mathbb{R}^n \oplus i\mathbb{R}^n, \quad \varepsilon \in \mathbb{R}$$

- Find a second map that leaves $\tilde{\Delta}(\varepsilon)$ invariant

$$\varpi : \tilde{\Delta}(\varepsilon) \rightarrow \tilde{\Delta}(\varepsilon), \quad \tilde{\alpha} \mapsto \omega \tilde{\alpha}$$

- (i) $\varpi : \tilde{\alpha} = \mu_1 \alpha_1 + \mu_2 \alpha_2 \mapsto \mu_1^* \omega \alpha_1 + \mu_2^* \omega \alpha_2$ for $\mu_1, \mu_2 \in \mathbb{C}$
- (ii) $\varpi \circ \varpi = \mathbb{I}$

Make the following assumptions

(i) ω decomposes as

$$\omega = \tau \hat{\omega} = \hat{\omega} \tau$$

with $\hat{\omega} \in \mathcal{W}$, $\hat{\omega}^2 = \mathbb{I}$ and complex conjugation τ

(ii) there are at least two different ω_i with $i = 1, 2, \dots$

(iii) there is a similarity transformation

$$\omega_i := \theta_\varepsilon \hat{\omega}_i \theta_\varepsilon^{-1} = \tau \hat{\omega}_i, \quad \text{for } i = 1, \dots, \kappa \geq 2$$

Make the following assumptions

(i) ω decomposes as

$$\omega = \tau \hat{\omega} = \hat{\omega} \tau$$

with $\hat{\omega} \in \mathcal{W}$, $\hat{\omega}^2 = \mathbb{I}$ and complex conjugation τ

(ii) there are at least two different ω_i with $i = 1, 2, \dots$

(iii) there is a similarity transformation

$$\omega_i := \theta_\varepsilon \hat{\omega}_i \theta_\varepsilon^{-1} = \tau \hat{\omega}_i, \quad \text{for } i = 1, \dots, \kappa \geq 2$$

(iv) θ_ε is an isometry for the inner products on $\tilde{\Delta}(\varepsilon)$ therefore

$$\theta_\varepsilon^* = \theta_\varepsilon^{-1} \quad \text{and} \quad \det \theta_\varepsilon = \pm 1$$

Make the following assumptions

(i) ω decomposes as

$$\omega = \tau \hat{\omega} = \hat{\omega} \tau$$

with $\hat{\omega} \in \mathcal{W}$, $\hat{\omega}^2 = \mathbb{I}$ and complex conjugation τ

(ii) there are at least two different ω_j with $j = 1, 2, \dots$

(iii) there is a similarity transformation

$$\omega_j := \theta_\varepsilon \hat{\omega}_j \theta_\varepsilon^{-1} = \tau \hat{\omega}_j, \quad \text{for } j = 1, \dots, \kappa \geq 2$$

(iv) θ_ε is an isometry for the inner products on $\tilde{\Delta}(\varepsilon)$ therefore

$$\theta_\varepsilon^* = \theta_\varepsilon^{-1} \quad \text{and} \quad \det \theta_\varepsilon = \pm 1$$

(v) in the limit $\varepsilon \rightarrow 0$ we recover the undeformed case

$$\lim_{\varepsilon \rightarrow 0} \theta_\varepsilon = \mathbb{I}$$

Make the following assumptions

(i) ω decomposes as

$$\omega = \tau \hat{\omega} = \hat{\omega} \tau$$

with $\hat{\omega} \in \mathcal{W}$, $\hat{\omega}^2 = \mathbb{I}$ and complex conjugation τ

(ii) there are at least two different ω_j with $j = 1, 2, \dots$

(iii) there is a similarity transformation

$$\omega_j := \theta_\varepsilon \hat{\omega}_j \theta_\varepsilon^{-1} = \tau \hat{\omega}_j, \quad \text{for } j = 1, \dots, \kappa \geq 2$$

(iv) θ_ε is an isometry for the inner products on $\tilde{\Delta}(\varepsilon)$ therefore

$$\theta_\varepsilon^* = \theta_\varepsilon^{-1} \quad \text{and} \quad \det \theta_\varepsilon = \pm 1$$

(v) in the limit $\varepsilon \rightarrow 0$ we recover the undeformed case

$$\lim_{\varepsilon \rightarrow 0} \theta_\varepsilon = \mathbb{I}$$

Make the following assumptions

(i) ω decomposes as

$$\omega = \tau \hat{\omega} = \hat{\omega} \tau$$

with $\hat{\omega} \in \mathcal{W}$, $\hat{\omega}^2 = \mathbb{I}$ and complex conjugation τ

(ii) there are at least two different ω_i with $i = 1, 2, \dots$

(iii) there is a similarity transformation

$$\omega_i := \theta_\varepsilon \hat{\omega}_i \theta_\varepsilon^{-1} = \tau \hat{\omega}_i, \quad \text{for } i = 1, \dots, \kappa \geq 2$$

(iv) θ_ε is an isometry for the inner products on $\tilde{\Delta}(\varepsilon)$ therefore

$$\theta_\varepsilon^* = \theta_\varepsilon^{-1} \quad \text{and} \quad \det \theta_\varepsilon = \pm 1$$

(v) in the limit $\varepsilon \rightarrow 0$ we recover the undeformed case

$$\lim_{\varepsilon \rightarrow 0} \theta_\varepsilon = \mathbb{I}$$

$\tilde{\Delta}(\varepsilon)$ for A_{4n-1} -subseries

closed solution

$$\theta_\varepsilon = r_0 \mathbb{I} + r_{2n} \sigma^{2n} + \imath r_n (\sigma^n - \sigma^{-n}),$$

- with $r_{2n} = 1 - r_0$, $r_n = \pm \sqrt{r_0^2 - r_0}$

- useful choice $r_0 = \cosh \varepsilon$

 $\tilde{\Delta}(\varepsilon)$ for E_6

$$\theta_\varepsilon = \begin{pmatrix} r_0 & -2\imath r_2 & 0 & -2\imath r_2 & -2\imath r_2 & -\imath r_2 \\ 2\imath r_2 & r_0 + \imath r_2 & 2\imath r_2 & 2\imath r_2 & 2\imath r_2 & 2\imath r_2 \\ 0 & 2\imath r_2 & r_0 + 2\imath r_2 & 4\imath r_2 & 3\imath r_2 & 2\imath r_2 \\ -2\imath r_2 & -2\imath r_2 & -4\imath r_2 & r_0 - 5\imath r_2 & -4\imath r_2 & -2\imath r_2 \\ 2\imath r_2 & 2\imath r_2 & 3\imath r_2 & 4\imath r_2 & r_0 + 2\imath r_2 & 0 \\ -\imath r_2 & -2\imath r_2 & -2\imath r_2 & -2\imath r_2 & 0 & r_0 \end{pmatrix}$$

$$r_2 = \pm 1/\sqrt{3} \sqrt{r_0^2 - 1}, \quad r_0 = \cosh \varepsilon$$

The generic case

- generalized Calogero Hamiltonian (undeformed)

$$\mathcal{H}_C(p, q) = \frac{p^2}{2} + \frac{\omega^2}{4} \sum_{\alpha \in \Delta^+} (\alpha \cdot q)^2 + \sum_{\alpha \in \Delta^+} \frac{g_\alpha}{(\alpha \cdot q)^2},$$

- define the variables

$$z := \prod_{\alpha \in \Delta^+} (\alpha \cdot q) \quad \text{and} \quad r^2 := \frac{1}{\hat{h}t_\ell} \sum_{\alpha \in \Delta^+} (\alpha \cdot q)^2,$$

$\hat{h} \equiv$ dual Coxeter number, $t_\ell \equiv \ell$ -th symmetrizer of l

- Ansatz:

$$\psi(q) \rightarrow \psi(z, r) = z^{\kappa+1/2} \varphi(r)$$

\Rightarrow solution for $\kappa = 1/2\sqrt{1+4g}$.

$$\varphi_n(r) = c_n \exp\left(-\sqrt{\frac{\hat{h}t_\ell}{2}} \frac{\omega}{2} r^2\right) L_n^a\left(\sqrt{\frac{\hat{h}t_\ell}{2}} \omega r^2\right).$$

$L_n^a(x) \equiv$ Laguerre polynomial, $a = (2 + h + h\sqrt{1+4g})l/4 - 1$

The generic case

- antilinearly deformed Calogero Hamiltonian

$$\mathcal{H}_{adC}(p, q) = \frac{p^2}{2} + \frac{\omega^2}{4} \sum_{\tilde{\alpha} \in \tilde{\Delta}^+} (\tilde{\alpha} \cdot q)^2 + \sum_{\tilde{\alpha} \in \Delta^+} \frac{g_{\tilde{\alpha}}}{(\tilde{\alpha} \cdot q)^2}$$

- define the variables

$$\tilde{z} := \prod_{\tilde{\alpha} \in \tilde{\Delta}^+} (\tilde{\alpha} \cdot q) \quad \text{and} \quad \tilde{r}^2 := \frac{1}{\hat{h}t_\ell} \sum_{\tilde{\alpha} \in \tilde{\Delta}^+} (\tilde{\alpha} \cdot q)^2$$

- Ansatz

$$\psi(q) \rightarrow \psi(\tilde{z}, \tilde{r}) = \tilde{z}^s \varphi(\tilde{r})$$

The generic case

- antilinearly deformed Calogero Hamiltonian

$$\mathcal{H}_{adC}(p, q) = \frac{p^2}{2} + \frac{\omega^2}{4} \sum_{\tilde{\alpha} \in \tilde{\Delta}^+} (\tilde{\alpha} \cdot q)^2 + \sum_{\tilde{\alpha} \in \Delta^+} \frac{g_{\tilde{\alpha}}}{(\tilde{\alpha} \cdot q)^2}$$

- define the variables

$$\tilde{z} := \prod_{\tilde{\alpha} \in \tilde{\Delta}^+} (\tilde{\alpha} \cdot q) \quad \text{and} \quad \tilde{r}^2 := \frac{1}{\hat{h}_{\ell}} \sum_{\tilde{\alpha} \in \tilde{\Delta}^+} (\tilde{\alpha} \cdot q)^2$$

- Ansatz

$$\psi(q) \rightarrow \psi(\tilde{z}, \tilde{r}) = \tilde{z}^s \varphi(\tilde{r})$$

when identities still hold \Rightarrow

$$\psi(q) = \psi(\tilde{z}, r) = \tilde{z}^s \varphi_n(r)$$

eigenenergies with different constraints (only performed for ground state)

Deformed A_3 -models

- potential from deformed Coxeter group factors

$$\alpha_1 = \{1, -1, 0, 0\}, \alpha_2 = \{0, 1, -1, 0\}, \alpha_3 = \{0, 0, 1, -1\}$$

$$\tilde{\alpha}_1 \cdot q = q_{43} + \cosh \varepsilon (q_{12} + q_{34}) - i\sqrt{2} \cosh \varepsilon \sinh \frac{\varepsilon}{2} (q_{13} + q_{24})$$

$$\tilde{\alpha}_2 \cdot q = q_{23} (2 \cosh \varepsilon - 1) + i2\sqrt{2} \cosh \varepsilon \sinh \frac{\varepsilon}{2} q_{14}$$

$$\tilde{\alpha}_3 \cdot q = q_{21} + \cosh \varepsilon (q_{12} + q_{34}) - i\sqrt{2} \cosh \varepsilon \sinh \frac{\varepsilon}{2} (q_{13} + q_{24})$$

$$\tilde{\alpha}_4 \cdot q = q_{42} + \cosh \varepsilon (q_{13} + q_{24}) + i\sqrt{2} \cosh \varepsilon \sinh \frac{\varepsilon}{2} (q_{12} + q_{34})$$

$$\tilde{\alpha}_5 \cdot q = q_{31} + \cosh \varepsilon (q_{13} + q_{24}) + i\sqrt{2} \cosh \varepsilon \sinh \frac{\varepsilon}{2} (q_{12} + q_{34})$$

$$\tilde{\alpha}_6 \cdot q = q_{14} (2 \cosh \varepsilon - 1) - i\sqrt{2} \cosh \varepsilon \sinh \frac{\varepsilon}{2} q_{23}$$

notation $q_{ij} = q_i - q_j$,

Deformed A_3 -models

- potential from deformed Coxeter group factors

$$\alpha_1 = \{1, -1, 0, 0\}, \alpha_2 = \{0, 1, -1, 0\}, \alpha_3 = \{0, 0, 1, -1\}$$

$$\tilde{\alpha}_1 \cdot q = q_{43} + \cosh \varepsilon (q_{12} + q_{34}) - i\sqrt{2} \cosh \varepsilon \sinh \frac{\varepsilon}{2} (q_{13} + q_{24})$$

$$\tilde{\alpha}_2 \cdot q = q_{23} (2 \cosh \varepsilon - 1) + i2\sqrt{2} \cosh \varepsilon \sinh \frac{\varepsilon}{2} q_{14}$$

$$\tilde{\alpha}_3 \cdot q = q_{21} + \cosh \varepsilon (q_{12} + q_{34}) - i\sqrt{2} \cosh \varepsilon \sinh \frac{\varepsilon}{2} (q_{13} + q_{24})$$

$$\tilde{\alpha}_4 \cdot q = q_{42} + \cosh \varepsilon (q_{13} + q_{24}) + i\sqrt{2} \cosh \varepsilon \sinh \frac{\varepsilon}{2} (q_{12} + q_{34})$$

$$\tilde{\alpha}_5 \cdot q = q_{31} + \cosh \varepsilon (q_{13} + q_{24}) + i\sqrt{2} \cosh \varepsilon \sinh \frac{\varepsilon}{2} (q_{12} + q_{34})$$

$$\tilde{\alpha}_6 \cdot q = q_{14} (2 \cosh \varepsilon - 1) - i\sqrt{2} \cosh \varepsilon \sinh \frac{\varepsilon}{2} q_{23}$$

notation $q_{ij} = q_i - q_j$, **No longer singular for $q_{ij} = 0$**

- \mathcal{PT} -symmetry for $\tilde{\alpha}$

$$\sigma_-^{\mathcal{E}} : \tilde{\alpha}_1 \rightarrow -\tilde{\alpha}_1, \tilde{\alpha}_2 \rightarrow \tilde{\alpha}_6, \tilde{\alpha}_3 \rightarrow -\tilde{\alpha}_3, \tilde{\alpha}_4 \rightarrow \tilde{\alpha}_5, \tilde{\alpha}_5 \rightarrow \tilde{\alpha}_4, \tilde{\alpha}_6 \rightarrow \tilde{\alpha}_2$$

$$\sigma_+^{\mathcal{E}} : \tilde{\alpha}_1 \rightarrow \tilde{\alpha}_4, \tilde{\alpha}_2 \rightarrow -\tilde{\alpha}_2, \tilde{\alpha}_3 \rightarrow \tilde{\alpha}_5, \tilde{\alpha}_4 \rightarrow \tilde{\alpha}_1, \tilde{\alpha}_5 \rightarrow \tilde{\alpha}_3, \tilde{\alpha}_6 \rightarrow \tilde{\alpha}_6$$

- \mathcal{PT} -symmetry in dual space

$$\sigma_-^{\mathcal{E}} : q_1 \rightarrow q_2, q_2 \rightarrow q_1, q_3 \rightarrow q_4, q_4 \rightarrow q_3, \iota \rightarrow -\iota$$

$$\sigma_+^{\mathcal{E}} : q_1 \rightarrow q_1, q_2 \rightarrow q_3, q_3 \rightarrow q_2, q_4 \rightarrow q_4, \iota \rightarrow -\iota$$

- \mathcal{PT} -symmetry for $\tilde{\alpha}$

$$\sigma_-^\varepsilon : \tilde{\alpha}_1 \rightarrow -\tilde{\alpha}_1, \tilde{\alpha}_2 \rightarrow \tilde{\alpha}_6, \tilde{\alpha}_3 \rightarrow -\tilde{\alpha}_3, \tilde{\alpha}_4 \rightarrow \tilde{\alpha}_5, \tilde{\alpha}_5 \rightarrow \tilde{\alpha}_4, \tilde{\alpha}_6 \rightarrow \tilde{\alpha}_2,$$

$$\sigma_+^\varepsilon : \tilde{\alpha}_1 \rightarrow \tilde{\alpha}_4, \tilde{\alpha}_2 \rightarrow -\tilde{\alpha}_2, \tilde{\alpha}_3 \rightarrow \tilde{\alpha}_5, \tilde{\alpha}_4 \rightarrow \tilde{\alpha}_1, \tilde{\alpha}_5 \rightarrow \tilde{\alpha}_3, \tilde{\alpha}_6 \rightarrow \tilde{\alpha}_6$$

- \mathcal{PT} -symmetry in dual space

$$\sigma_-^\varepsilon : q_1 \rightarrow q_2, q_2 \rightarrow q_1, q_3 \rightarrow q_4, q_4 \rightarrow q_3, \iota \rightarrow -\iota$$

$$\sigma_+^\varepsilon : q_1 \rightarrow q_1, q_2 \rightarrow q_3, q_3 \rightarrow q_2, q_4 \rightarrow q_4, \iota \rightarrow -\iota$$

\Rightarrow

$$\sigma_-^\varepsilon \tilde{z}(q_1, q_2, q_3, q_4) = \tilde{z}^*(q_2, q_1, q_4, q_3) = \tilde{z}(q_1, q_2, q_3, q_4)$$

$$\sigma_+^\varepsilon \tilde{z}(q_1, q_2, q_3, q_4) = \tilde{z}^*(q_1, q_3, q_2, q_4) = -\tilde{z}(q_1, q_2, q_3, q_4)$$

$$\psi(q_1, q_2, q_3, q_4) = e^{i\pi S} \psi(q_2, q_4, q_1, q_3).$$

Anyonic exchange factors in the 4-particle scattering process

$$\begin{array}{cccc}
 w & x & y & z \\
 \bullet & \bullet & \bullet & \bullet \\
 q_1 & q_2 & q_3 & q_4
 \end{array}
 = e^{2\pi S}
 \begin{array}{cccc}
 w & x & y & z \\
 \bullet & \bullet & \bullet & \bullet \\
 q_2 & q_4 & q_1 & q_3
 \end{array}$$

Anyonic exchange factors in the 4-particle scattering process

$$\begin{array}{cccc}
 w & x & y & z \\
 \bullet & \bullet & \bullet & \bullet \\
 q_1 & q_2 & q_3 & q_4
 \end{array}
 = e^{2\pi S}
 \begin{array}{cccc}
 w & x & y & z \\
 \bullet & \bullet & \bullet & \bullet \\
 q_2 & q_4 & q_1 & q_3
 \end{array}$$

$$\begin{array}{ccc}
 x & y & z \\
 \bullet & \bullet & \bullet \\
 q_1 & q_2 = q_3 & q_4
 \end{array}
 = e^{2\pi S}
 \begin{array}{ccc}
 x & y & z \\
 \bullet & \bullet & \bullet \\
 q_2 & q_1 = q_4 & q_3
 \end{array}$$

Anyonic exchange factors in the 4-particle scattering process

$$\begin{array}{cccc}
 w & x & y & z \\
 \bullet & \bullet & \bullet & \bullet \\
 q_1 & q_2 & q_3 & q_4
 \end{array}
 = e^{2\pi S}
 \begin{array}{cccc}
 w & x & y & z \\
 \bullet & \bullet & \bullet & \bullet \\
 q_2 & q_4 & q_1 & q_3
 \end{array}$$

$$\begin{array}{ccc}
 x & y & z \\
 \bullet & \bullet & \bullet \\
 q_1 & q_2 = q_3 & q_4
 \end{array}
 = e^{2\pi S}
 \begin{array}{ccc}
 x & y & z \\
 \bullet & \bullet & \bullet \\
 q_2 & q_1 = q_4 & q_3
 \end{array}$$

$$\begin{array}{cc}
 x & y \\
 \bullet & \bullet \\
 q_1 = q_2 & q_3 = q_4
 \end{array}
 = e^{2\pi S}
 \begin{array}{cc}
 x & y \\
 \bullet & \bullet \\
 q_1 = q_3 & q_2 = q_4
 \end{array}$$

Find Hermitian counterpart h , Dyson map η and metric ρ :

$$h = \eta H \eta^{-1} = h^\dagger = (\eta^{-1})^\dagger H^\dagger \eta^\dagger \Leftrightarrow H^\dagger \rho = \rho H \quad \text{with } \rho = \eta^\dagger \eta$$

Some B_ℓ -models correspond to complex rotations

$$\begin{pmatrix} \tilde{z}_i \\ \tilde{z}_j \end{pmatrix} = R_{ij} \begin{pmatrix} z_i \\ z_j \end{pmatrix} = \eta_{ij} \begin{pmatrix} z_i \\ z_j \end{pmatrix} \eta_{ij}^{-1}, \quad \text{for } z \in \{x, p\}, \quad \eta_{ij} = e^{\varepsilon(x_i p_j - x_j p_i)}$$

For instance for:

$$\theta_\varepsilon^* = \begin{pmatrix} R & & & \\ & R & & 0 \\ & & R & \\ & 0 & & \ddots \\ & & & & 1 \end{pmatrix} \quad \text{with } R = \begin{pmatrix} \cosh \varepsilon & i \sinh \varepsilon \\ -i \sinh \varepsilon & \cosh \varepsilon \end{pmatrix}$$

we have

$$\mathcal{H}_0(p, x) = \eta \mathcal{H}_\varepsilon(p, x) \eta^{-1}$$

with

$$\eta = \eta_{12}^{-1} \eta_{34}^{-1} \eta_{56}^{-1} \cdots \eta_{(\ell-2)(\ell-1)}^{-1}$$

For B_5

$$\theta_\varepsilon^* = \begin{pmatrix} r_0 & -i\vartheta & i\vartheta & 1-r_0 & 0 \\ i\vartheta & r_0 & 1-r_0 & -i\vartheta & 0 \\ -i\vartheta & 1-r_0 & r_0 & i\vartheta & 0 \\ 1-r_0 & i\vartheta & -i\vartheta & r_0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

we find

$$\tilde{X} = \theta_\varepsilon^* X = R_{24}^{-1} R_{13} R_{34} R_{12}^{-1} X = \eta X \eta^{-1}, \quad \text{with } \eta = \eta_{24}^{-1} \eta_{13} \eta_{34} \eta_{12}^{-1}.$$

In general this is an open problem.

General deformation prescription:

\mathcal{PT} -anti-symmetric quantities:

$$\mathcal{PT} : \phi(x, t) \mapsto -\phi(x, t) \quad \Rightarrow \quad \delta_\varepsilon : \phi(x, t) \mapsto -i[i\phi(x, t)]^\varepsilon$$

Two possibilities for the KdV Hamiltonian

$$\delta_\varepsilon^+ : u_x \mapsto u_{x,\varepsilon} := -i(iu_x)^\varepsilon \quad \text{or} \quad \delta_\varepsilon^- : u \mapsto u_\varepsilon := -i(iu)^\varepsilon,$$



such that

$$\mathcal{H}_\varepsilon^+ = -\frac{\beta}{6}u^3 - \frac{\gamma}{1+\varepsilon}(iu_x)^{\varepsilon+1} \quad \mathcal{H}_\varepsilon^- = \frac{\beta}{(1+\varepsilon)(2+\varepsilon)}(iu)^{\varepsilon+2} + \frac{\gamma}{2}u_x^2$$

with equations of motion

$$u_t + \beta uu_x + \gamma u_{xxx,\varepsilon} = 0 \quad u_t + i\beta u_\varepsilon u_x + \gamma u_{xxx} = 0$$



Broken \mathcal{PT} -symmetric rational solutions for $\mathcal{H}_{1/3}^+$



Different Riemann sheets for $A = (1 - i)/4$, $c = 1$, $\beta = 2 + 2i$
and $\gamma = 3$

(a) $u^{(1)}$

(b) $u^{(2)}$



The $\mathcal{H}_\varepsilon^+$ -models **\mathcal{PT} -symmetric trigonometric/hyperbolic solutions**

$A = 4, B = 2, c = 1, \beta = 2$ and $\gamma = 3$

(a) $\mathcal{H}_{-1/2}^+$

(b) $\mathcal{H}_{-2/3}^+$



Broken \mathcal{PT} -symmetric trigonometric solutions for $\mathcal{H}_{-1/2}^+$



(a) Spontaneously broken \mathcal{PT} -symmetry with $A = 4 + i$,
 $B = 2 - 2i$, $c = 1$, $\beta = 3/10$ and $\gamma = 3$

(b) broken \mathcal{PT} -symmetry with $A = 4$, $B = 2$, $c = 1$, $\beta = 3/10$
 and $\gamma = 3 + i$

Elliptic solutions for $\mathcal{H}_{-1/2}^+$:



(a) \mathcal{PT} -symmetric with $A = 1$, $B = 3$, $C = 6$, $\beta = 3/10$, $\gamma = -3$ and $c = 1$

(b) spontaneously broken \mathcal{PT} -symmetry with $A = 1 + i$, $B = 3 - i$, $C = 6$, $\beta = 3/10$, $\gamma = -3$ and $c = 1$

The $\mathcal{H}_\varepsilon^-$ -models

Integrating twice gives now:

$$u_\zeta^2 = \frac{2}{\gamma} \left(\kappa_2 + \kappa_1 u + \frac{c}{2} u^2 - \beta \frac{i^\varepsilon}{(1+\varepsilon)(2+\varepsilon)} u^{2+\varepsilon} \right) =: \lambda Q(u)$$



where

$$\lambda = -\frac{2\beta i^\varepsilon}{\gamma(1+\varepsilon)(2+\varepsilon)}$$

For $\kappa_1 = \kappa_2 = 0$

$$u(\zeta) = \left(\frac{c(\varepsilon+1)(\varepsilon+2)}{i^\varepsilon \beta \left[\cosh \left(\frac{\sqrt{c\varepsilon}(\zeta-\zeta_0)}{\sqrt{\gamma}} \right) + 1 \right]} \right)^{1/\varepsilon}$$

Broken \mathcal{PT} -symmetric solution for \mathcal{H}_4^- :



(a) star node at the origin for $c = 1$, $\beta = 2 + i3$, $\gamma = 1$ and $B = (15/2 + i3)^{1/4}$

(b) centre at the origin for $c = 1$, $\beta = 2 + i3$, $\gamma = -1$ and $B = (30/13 - i45/13)^{1/4}$

Reduction of the \mathcal{H}_2^- -model

$$\mathcal{H}_2^-[u] = \frac{\beta}{12}u^4 + \frac{\gamma}{2}u_x^2$$

Twice integrated equation of motion:

$$u_\zeta^2 = \frac{2}{\gamma} \left(\kappa_2 + \kappa_1 u + \frac{c}{2}u^2 + \beta \frac{1}{12}u^4 \right) =: \lambda Q(u)$$

Reduction $u \rightarrow x, \zeta \rightarrow t$

$$\kappa_1 = -\gamma\tau, \quad \kappa_2 = \gamma E_x, \quad \beta = -3\gamma g \quad \text{and} \quad c = -\gamma\omega^2$$

Quartic harmonic oscillator of the form

$$H = E_x = \frac{1}{2}p^2 + \tau x + \frac{\omega^2}{2}x^2 + \frac{g}{4}x^4$$

Boundary cond.: $\kappa_1 = \tau = 0, \lim_{\zeta \rightarrow \infty} u(\zeta) = 0, \lim_{\zeta \rightarrow \infty} u_x(\zeta) = \sqrt{2E_x}$

[A.G. Anderson, C. Bender, U. Morone, arXiv:1102.4822]

Reduction of the \mathcal{H}_2^- -model

$$\mathcal{H}_2^-[u] = \frac{\beta}{12}u^4 + \frac{\gamma}{2}u_x^2$$

Twice integrated equation of motion:

$$u_\zeta^2 = \frac{2}{\gamma} \left(\kappa_2 + \kappa_1 u + \frac{c}{2}u^2 + \beta \frac{1}{12}u^4 \right) =: \lambda Q(u)$$

Reduction $u \rightarrow x, \zeta \rightarrow t$

$$\kappa_1 = -\gamma\tau, \quad \kappa_2 = \gamma E_x, \quad \beta = -3\gamma g \quad \text{and} \quad c = -\gamma\omega^2$$

Quartic harmonic oscillator of the form

$$H = E_x = \frac{1}{2}p^2 + \tau x + \frac{\omega^2}{2}x^2 + \frac{g}{4}x^4$$

Boundary cond.: $\kappa_1 = \tau = 0, \lim_{\zeta \rightarrow \infty} u(\zeta) = 0, \lim_{\zeta \rightarrow \infty} u_x(\zeta) = \sqrt{2E_x}$

[A.G. Anderson, C. Bender, U. Morone, arXiv:1102.4822]

Note: $E_x \neq E_u(a)$

Ito type systems and its deformations

Coupled nonlinear system

$$\begin{aligned}u_t + \alpha v v_x + \beta u u_x + \gamma u_{xxx} &= 0, & \alpha, \beta, \gamma \in \mathbb{C}, \\v_t + \delta(uv)_x + \phi v_{xxx} &= 0, & \delta, \phi \in \mathbb{C}\end{aligned}$$

Hamiltonian for $\delta = \alpha$

$$\mathcal{H}_I = -\frac{\alpha}{2} uv^2 - \frac{\beta}{6} u^3 + \frac{\gamma}{2} u_x^2 + \frac{\phi}{2} v_x^2$$

\mathcal{PT} -symmetries:

$$\mathcal{PT}_{++} : x \mapsto -x, t \mapsto -t, i \mapsto -i, u \mapsto u, v \mapsto v \quad \text{for } \alpha, \beta, \gamma, \phi \in \mathbb{R}$$

$$\mathcal{PT}_{+-} : x \mapsto -x, t \mapsto -t, i \mapsto -i, u \mapsto u, v \mapsto -v \quad \text{for } \alpha, \beta, \gamma, \phi \in \mathbb{R}$$

$$\mathcal{PT}_{-+} : x \mapsto -x, t \mapsto -t, i \mapsto -i, u \mapsto -u, v \mapsto v \quad \text{for } i\alpha, i\beta, \gamma, \phi \in \mathbb{R}$$

$$\mathcal{PT}_{--} : x \mapsto -x, t \mapsto -t, i \mapsto -i, u \mapsto -u, v \mapsto -v \quad \text{for } i\alpha, i\beta, \gamma, \phi \in \mathbb{R}$$

Deformed models

$$\mathcal{H}_{\varepsilon,\mu}^{++} = -\frac{\alpha}{2}uv^2 - \frac{\beta}{6}u^3 - \frac{\gamma}{1+\varepsilon}(iu_x)^{\varepsilon+1} - \frac{\phi}{1+\mu}(iv_x)^{\mu+1}$$

$$\mathcal{H}_{\varepsilon,\mu}^{+-} = \frac{\alpha}{1+\mu}u(iv)^{\mu+1} - \frac{\beta}{6}u^3 - \frac{\gamma}{1+\varepsilon}(iu_x)^{\varepsilon+1} + \frac{\phi}{2}v_x^2$$

$$\mathcal{H}_{\varepsilon,\mu}^{-+} = -\frac{\alpha}{2}uv^2 - \frac{i\beta}{(1+\varepsilon)(2+\varepsilon)}(iu)^{2+\varepsilon} + \frac{\gamma}{2}u_x^2 - \frac{\phi}{1+\mu}(iv_x)^{\mu+1}$$

$$\mathcal{H}_{\varepsilon,\mu}^{--} = \frac{\alpha}{1+\mu}u(iv)^{\mu+1} - \frac{i\beta}{(1+\varepsilon)(2+\varepsilon)}(iu)^{2+\varepsilon} + \frac{\gamma}{2}u_x^2 + \frac{\phi}{2}v_x^2$$

with equations of motion

$$\begin{aligned} u_t + \alpha vv_x + \beta uu_x + \gamma u_{xxx,\varepsilon} &= 0, & u_t + \alpha v_\mu v_x + \beta uu_x + \gamma u_{xxx,\varepsilon} &= 0, \\ v_t + \alpha (uv)_x + \phi v_{xxx,\mu} &= 0, & v_t + \alpha (uv_\mu)_x + \phi v_{xxx} &= 0, \end{aligned}$$

$$\begin{aligned} u_t + \alpha vv_x + \beta u_\varepsilon u_x + \gamma u_{xxx} &= 0, & u_t + \alpha v_\mu v_x + \beta u_\varepsilon u_x + \gamma u_{xxx} &= 0, \\ v_t + \alpha (uv)_x + \phi v_{xxx,\mu} &= 0, & v_t + \alpha (uv_\mu)_x + \phi v_{xxx} &= 0. \end{aligned}$$

